

The Soviet Far East; Economic and Geographic Features (Cont.)	245
Ch. I. Physical and Geographic Features and Natural Resources	9
Topography	9
Minerals	17
Lakes and rivers	22
Climate	28
Permafrost	32
Soils	33
Flora	36
Fauna	45

Card 3/5

The Soviet Far East; Economic and Geographic Features (Cont.)	245
Ch. II. Stages of Colonization	48
Ch. III. Population	67
Ch. IV. Economy	77
Industries	78
Agriculture and animal husbandry	102
Transportation and interregional communications	121
Ch. V. Features of Individual Krays and Oblasts	134
Khabarovskiy Kray	134
Primorskiy Kray	154

Card 4/5

The Soviet Far East; Economic and Geographic Features (Cont.)	245
Amurskaya Oblast	176
Sakhalinskaya Oblast	192
Kamchatskaya Oblast	211
Magadanskaya Oblast	221
Bibliography	235
Appendix	237

AVAILABLE: Library of Congress

GC/bmd
May 28, 1958

Card 5/5

KLYUCHEROV, A.P.; KONDRAT'YEV, S.N.; Prinimali uchastiye: GUSAROV, F.V.;
UDOVENKO, V.G.; PETROV, G.A.; BURKSER, V.Ye.; SHMONIN, I.A.;
KUDRIN, Ye.A.; GALAKHMATOV, S.N.; ZIMINA, L.P.; SHISHARIN, B.N.;
KONDYURINA, R.V.; BURMISTROV, K.A.; SHIRNIN, I.A.; SIMONENKO, F.N.;
GORSHILOV, Yu.V.; KOLPAKOV, B.V.; GUSAROV, A.K.; BOLOTOV, P.G.

Heat insulation of open-hearth furnace crowns. Metallurg 5 no.11:
14-17 N '60. (MIRA 13:10)

1. Nizhe-Tagil'skiy metallurgicheskiy kombinat.
(Open-hearth furnaces--Design and construction)
(Insulation (Heat))

KONDRAT'YEV, S.N.; KLYUCHEROV, A.P.; UDOVENKO, V.G.; SHIRNIN, I.A.;
VYDRINA, Zh.A.

Rapid methods of repair and the fritting of new hearth bottoms.
Metallurg 6 no.9-10-13 S '61. (MIRA 14:9)

1. Nizhne-Tagil'skiy metallurgicheskiy kombinat.
(Open-hearth furnaces--Maintenance and repair)

BARANOV, V.M.; DONSKOY, S.A.; TORSHILOV, Yn.V.; TRET'YAKOV, M.A.; UDOLYENKO,
V.G.; FREYDENZON, Ye.Z.

Blowing of cast iron in high-capacity converters. Metallurg 10 no.9:
15-18 S '65. (MIRA 18:9)

1. Nizhne-Tagil'skiy metallurgicheskiy kombinat.

ALAMPIYEV, P.M.; OMAROVSKIY, A.G.; UDOVENKO, V.A.

New features of the literature on the economic regions of the
U.S.S.R. Izv. AN SSSR. Ser. geog. no.4:147-151 J1-Ag '63.

(MIRA 16:8)

(Bibliography--Geography, Economic)

DUBROV, N.F.; KITAYEV, B.I.; KOKAREV, N.I.; UDOVENKO, V.G.; KONDRAT'YEV, S.N.;
ZATULOVSKAYA, Ye.Z.; KLYUCHEROV, A.P.

Review of the book by N.A.Vecher "Highly efficient operation of
open-hearth furnaces." Stal' 24 no.7:613-614 J1 '64.

(MIRA 18:1)

1. Ural'skiy nauchno-issledovatel'skiy institut chernykh metallov,
Ural'skiy politekhnicheskoy institut i Nizhne-Tagil'skiy metallurgi-
cheskiy kombinat.

ARNAUTOV, V.T.; BARANOV, V.M.; DONSKOY, S.A.; PASTUKHOV, A.I.; SMIRNOV, I.A.; TORSHILOV, Yu.V.; TRET'YAKOV, M.A.; UDOVENKO, V.G.; FREYDENZON, Ye.Z.; SHCHEKALEV, Yu.S.; Prinimali uchastiye: MAKAYEV, S.V.; KOMPANIYETS, G.M.; NAGOVITSYN, D.F.; NOVOLODSKIY, P.I.; VARSHAVSKIY, V.L.; KOROGODSKIY, V.G.; KLIBANOV, Ye.L.; MEDVEDEVSKIY, Yu.; TALANTSEVA, T.I.; DUBROV, N.F.; DZEMIAN, S.K.; TOPYCHKANOV, B.I.; CHARUSHNIKOV, O.A.; KHARITONOV, Yu.A.

Developing and mastering the technology of converting vanadium cast iron in oxygen-blown converters with a 100 ton (Mg) capacity. Stal' 25 no.6:50%-508 Je '65. (MIRA 18:6)

1. Nizhne-Tagil'skiy metallurgicheskiy kombinat (for Makayev, Kompaniyets, Nagovitsyn, Novolodskiy, Varshavskiy, Korogodskiy, Klibanov, Medvedevskikh, Talantseva). 2. Ural'skiy nauchno-issledovatel'skiy institut chenykh metallov (for Dubrov, Dzemyan, Topychkanov, Charushnikov, Kharitonov).

FRIDYDENZON, Yo.Z.; UDDEVENOV, V.G.; TOREBNICOV, Yo.V.; KOMPANIYEV, V.M.;
TRET'YAKOV, M.A.; BARANOV, V.M.; MAGOVITSIN, L.F.; LANSKOY, S.A.;
PASTUKHOV, A.I.

Mastering the operation of the oxygen-blown converter plant
of the Nizhniy Tagil metallurgical combine. Stat' 25 no.6:
534-537 Je '65. (MIRA 12:6)

1. Nizhne-Tagil'skiy metallurgicheskiy kombinat i Ural'skiy
nauchno-issledovatel'skiy institut Chernykh metallov.

VARLAMOV, V.S., kand.tekhn.nauk; IL'INA, A.I.; KUDRYASHOV, A.I., inzh.;
UDOVENKO, V.S., inzh.; KOGAN, G.A., inzh.

Continuous oxidation of paraffins under industrial conditions. Masl.-zhir.prom. 25 no.10:39-41 '59.

(MIRA 13:2)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut zhirov (for Varlamov, Il'ina). 2. Shebekinskiy kombinat sinteticheskikh zhirnykh kislot i zhirnykh spirtov (for Kudryashov, Udovenko, Kogan).

(Shebekino--Paraffins)

1ST AND 2ND ORDERS																										3RD AND 4TH ORDERS																									
SUBJECTS AND PROPERTIES INDEX																										SUBJECTS AND PROPERTIES INDEX																									
<p>Specific heats of saturated vapors at the boiling point. N. A. KOTASOVSKII AND V. V. UDOLYNSKIY. <i>J. Gen. Chem.</i> (U. S. S. R.) 1, 255-62 (1931). Formulas expressing latent heat of vaporization and C_p as functions of abs. temp. are given for 20 liquids (H_2O, NH_3, $CHCl_3$, CCl_4, $MeOH$, Me_2CO, $EtCO_2H$, CaH_2, $EtCl$, $EtBr$, Et_2O, Me_2CH, CH_3Me, CaH_2, Et_2O, CS_2, $EtOH$, $AcOMe$, CaH_2 and CaH_2). For the calcn of sp. heat of satd vapors the approx thermodynamic formula $C_p = C_v + (\lambda/T)$ is used, where C_v is mol. sp. heat of satd vapor, C_p is mol. sp. heat of the vapor at const. pressure, λ = mol. latent heat of vaporization of the liquid and T = abs. temp. This formula can be used to verify results found through the application of the exact thermodynamic formula: $C_p = C_v + (d\lambda/dT) \cdot (\lambda/T)$, where C_v = mol. sp. heat of the liquid under pressure of satd. vapor. In most cases (first 13 liquids) the agreement between the 2 formulas is satisfactory. A rule for detg. the sign of sp. heat of a satd. vapor was found and verified. This rule is $C_p (n - 10) > 0$, where n is number of atoms in the mol. Abs. temp. of inversion of sign of C_p was calcd. for 14 liquids as follows: H_2O, 770°; $MeOH$, 461°; $CHCl_3$, 355°; $AcOMe$, 363° and 463°; Me_2CO, 341°; $EtBr$, 332°; CaH_2, 328°; CS_2, 323°; $EtCO_2H$, 308°; CaH_2, 299°; Me_2CHCH_2Me, 297° and 922°; Et_2O, 292°; CaH_2, 107° and 462°; $EtCl$, 144°.</p>																																																			
<p>ASAC-SLA DETAILING LITERATURE CLASSIFICATION</p>																																																			

Specific Heat of Liquids. II. N. A. Kolosovskii and V. V. Udovenko. *J. Gen. Chem.* (U. S. S. R.) 4, 1027-33 (1934); cf. C. A. 30, 3649¹. C_p in cal. per g. mol. at 18-30°, and temp. coeff. of entropy ($\Delta S/\Delta T$), were detd. for a number of liquids. The following figures give C_p ($\Delta S/\Delta T$), and temp.: resp. Benzene 31.30, 0.1090, 14.76; C_6H_6 , 42.73, 19.62, 0.2032, 20.20; sabinene 0.1574, 28.87; limonene 30.21, 0.1047, 15.11; $C_{11}H_{18}$ 34.32, 0.2031, 23.84; CCl_4 40.03, 0.1354, 20.19; 34.77, 0.1149, 20.45; C_6H_5CHO 40.03, 0.1354, 20.19; methyl ethyl ketone 34.45, 0.1200, 21.81; lincolol 39.05 (0.3034, 19.05); pulegone 35.37, 0.1240, 21.14; ethyl formate 35.37, 0.1200, 21.82; ethyl butyrate 39.10; formate 35.37, 0.1200, 21.82; ethyl benzoate 37.84, 0.1078, 19.34; benzyl acetate 39.78, 0.2042, 19.51; ethyl benzoate 37.84, 0.1078, 19.34; ethyl 0.1955; methyl salicylate 39.46, 0.2014, 22.00; carbonate 50.28, 0.1710, 21.50; diethyl succinate 30.77, 0.2700, 19.42; ethyl acetoacetate 50.28, 0.2009, 24.39; *m*-nitrotoluene 48.28, 0.1600, 29.19; *p*-toluidine 50.09, 0.1636, 29.33; *m*-toluidine 51.84, 0.1712, 29.52; di-methylaniline 51.29, 0.1606, 29.15; diethyl *p*. A table 0.2171, 28.83; quinoline 46.13, 0.1625, 29.21. of other shows a comparison of these results with those of other investigators. S. L. Mado.

COMMON ELEMENTS										PROCESSES AND PROPERTIES INDEX										1ST AND 2ND ORDERS										3RD AND 4TH ORDERS									
<p>OVONENKO, V. V.</p>										<p>Viscosity of the system ketones organic acids. I. V. V. Udavskiy. <i>J. Gen. Chem.</i> (U. S. S. R.) 9, 1163-6 (1938).—The viscosity and d. were detd. for binary systems contg. acetone and $\text{C}_2\text{H}_5\text{CO}_2\text{H}$ (at 35, 45, 55°), $\text{CH}_3\text{CICO}_2\text{H}$ (at 25, 35, 50°), HCO_2H (at 25, 35, 45°), resp. AcOH (at 20, 30, 40°), $\text{C}_6\text{H}_5\text{CO}_2\text{H}$ (at 25, 35, 45°), resp. A chem. reaction occurs between acetone and the acids. studied: Gertrude Berend</p>																													
<p>ASB-SEA METALLURGICAL LITERATURE CLASSIFICATION</p>										<p>1ST AND 2ND ORDERS</p>										<p>3RD AND 4TH ORDERS</p>																			

1ST AND 2ND GROUPS		PROCESSING AND PROPERTIES INDEX		3RD AND 4TH GROUPS	
<p><i>BC</i></p> <p>Viscosity in the systems: ketones-organic acids. II, III. Y. V. ILYUSHKIN (J. Gen. Chem. Russ., 1939, 9, 1512-1514, 1564-1567; cf. A., 1939, I, 606).—II. The η curves of the systems COMeEt or $\text{COMePr}-\text{HCO}_2\text{H}$ and $-\text{AcOH}$, at 25°, 36°, and 45°, do not suggest compound formation. —III. Shrinkage taking place when COMe_2 is mixed with HCO_2H, AcOH, or PrCO_2H is max. for equimol. mixtures, and increases in the order given. The same applies to the systems $\text{AcOH}-\text{COMePr}$, $-\text{COMeEt}$, and $-\text{COMe}_2$. This suggests chemical reaction between the components, although the η-composition curves furnish no evidence of it.</p> <p>R. T.</p>					
<p>ASTM-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>					
1ST GROUP		2ND GROUP		3RD GROUP	
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100		1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100		1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100	

1ST AND 2ND ORDER										3RD AND 4TH ORDER									
PROCESSING AND PROPERTIES INDEX																			
BC										A-1									
<p>Viscosity in the systems ketones-organic acids. IV. V. V. UDOLYUKO and R. P. AINA-PTOVA. V. V. UDOLYUKO and S. I. VITSANVA (J. Gen. Chem. Russ., 1939, 9, 1796-1800, 1731-1733; cf. A., 1940, 1, 66).—IV. The composition-η curves of the systems HCO_2H-, AcOH-, or PrCO_2H-C_6H_5-COMePr, at 25°, are concave to the composition axis, whilst in absence of C_6H_5 they are convex. This is ascribed to decrease in association of the acids due to dilution of the system. The curves suggest compound formation between COMePr and the org. acids to a degree diminishing in the order given.</p> <p>V. Analogous results are obtained in the systems HCO_2H-, AcOH-, or PrCO_2H-COMe-, C_6H_5, at 25°.</p> <p>R. T.</p>																			
A.S.T.M. METALLURGICAL LITERATURE CLASSIFICATION																			
FROM SYNONYM										FROM DOMINANT									
SYNONYM										SYNONYM									

1. USANOVICH, M., SUMAROKOVA, T., UDOVENKO, V.

2. USSR (600)

"Electrical Conductivity, Viscosity, and Surface Tension of the System $H_2SO_4-HClO_4$,"
Zhur. Obshch., 9, No. 21, 1939. Lab. of Physical Chem., Central Asiatic State
Univ. Received 9 June 1939.

9. ~~████~~ Report U-1626, 11 Jan 1952.

1ST AND 2ND ORDERS										3RD AND 4TH ORDERS									
UDOVENKO, V.V.																			
PROCESS AND PROPERTIES INDEX																			
AC																			
F 1																			
<p>Surface tension of systems ketones-organic acids. V. V. UDOVENKO, E. V. SITSCHKOVA, and A. P. TOMOROV (J. Gen. Chem. Russ., 1939, 9, 2048-2054).--The surface tension-composition curves of the systems COMe_2- or COMePr-$\text{R-CO}_2\text{H}$ ($\text{R} = \text{H, Me, Pr}$) and COMeEt-AcOH, at 25°, 35°, and 45°, deviate positively or negatively from the additive rule; these deviations are ascribed to shrinkage or dilatation of the liquids after mixing, rather than to formation of compounds.</p> <p>R. T.</p>																			
ASR-31A METALLURGICAL LITERATURE CLASSIFICATION																			
FROM SYNTHESE										FROM ANALYSIS									
SUBGROUP										SUBGROUP									
SUBGROUP										SUBGROUP									

PROCESSES AND PROPERTIES INDEX

Cryoscopic study of the systems phenol-aniline, phenol-dimethylaniline and phenol-diethylaniline. V. V. Udevenko and M. I. Usnikovich. *J. Gen. Chem.* (U. S. S. R.) 10, 17-20(1940); cf. preceding abstr.—The mol. wts. were detd. from the changes in the f. ps. of PhNH₂, Ph-NMe₂, and PhNEt₂, and their mixts. with PhOH in CCl₄, by the method previously described. Similar procedure was used in the detn. of mol. wts. of PhNH₂, and its alkyl deriva. in PhOH. The inconclusive results indicate that the mol. wts. of the mixts. with PhOH in CCl₄ are higher than the calcd. values; this can be explained by the formation of chem. compds. in the systems. The mol. wts. of anilines in PhOH decrease with increasing concn., approaching the normal values
Chas. Blanc

ASS-SLA METALLURGICAL LITERATURE CLASSIFICATION

1ST AND 2ND GROUPS										3RD AND 4TH GROUPS									
PROCESSES AND PROPERTIES INDEX																			
<div style="position: relative;"> B A-1 <div style="margin-top: 50px;"> <p>(A) Viscosity in the ternary systems: phenol-benzene-aniline, -dimethylaniline, and -diethylaniline. V. V. UDovenko and A. P. TOROFV.</p> <p>(B) Cryoscopic study of the systems phenol-aniline, -dimethylaniline, and -diethylaniline. V. V. UDovenko and M. I. USANOVITICH (J. Gen. Chem. Russ., 1940, 10, 11-16, 17-20).—(A) The η-composition curves suggest formation of $\text{PhOH} \cdot \text{NH}_2\text{Ph}$, -NPhMe_2, and -NPhEt_2 compounds; the MAX. become less well-defined as the $[\text{C}_6\text{H}_5]$ rises.</p> <p>(B) The apparent mol. wt. of NH_2Ph and NPhMe_2 in C_6H_6 rises with increasing concn.; that of NPhEt_2 remains const. In PhOH the apparent mol. wt. falls with rising concn. The mean apparent mol. wt. of mixtures of PhOH and amines in C_6H_6 is > would follow from the additive rule, to a degree increasing with their concn.</p> <p style="text-align: right;">R. T.</p> </div> </div>																			
ASB-51A METALLURGICAL LITERATURE CLASSIFICATION																			
1ST GROUP										2ND GROUP									
SUBGROUP										SUBGROUP									

1ST AND 2ND ORDERS										3RD AND 4TH ORDERS									
PROPERTIES AND PROPERTIES INDEX																			
<p>CA</p> <p>Viscosity of the systems acetic acid-dimethylaniline and acetic acid-diethylaniline. V. V. Udravko. J. Gen. Chem. (U. S. S. R.) 18, 1923-5(1940). The η was detd. at 25, 45 and 65°. The mol. % of HOAc ranged up to 100. The η curves pass through a max. at all three temps. and the max. is shifted with increasing temp. to the side of HOAc; or to the less-viscous component. In dimethyl-aniline-HOAc the max. was 85 mol. % of HOAc and in diethylaniline-HOAc at 90 mol. % HOAc. The latter solns. were more viscous than the former. The solns. evolved heat when prepd. and in both systems the formation of chem. compds. was noted but their compn. was not detd.</p> <p>H. Z. Kamich</p>																			
<p>ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION</p> <p>VSOMI SYMBOLIC VSOMI MIX OMV GSE VSOMI BOMINY VSOMI GSE OMV ASI</p>																			

Determination of Molecular Weight as a method of physicochemical analysis. V. Y. Lidskanyo. *J. Gen. Chem.* (U. S. S. R.) 11, 270-84 (1941). Cryometric data, made with mixts. of 2 components $A + B$, lead to values of a "mol. wt." of the two-component solute, defined by and calcd. from $M = (1000K - M_0\Delta T)/\Delta\Delta T$, where ΔT is the observed lowering of the f. p.; Δ , the wt. of the mixed solute ($A + B$); K , M_0 , B_0 , the cryometric const., mol. wt. and wt. of the solvent, resp. For several mol. ratios $A:B$, the so-detd. M is plotted against the total concn. of ($A + B$) (in mol. per 100 g. solvent) in the soln. Next, plots of M against the mol. ratio $A:B$ are constructed, (isocoon. lines). Measurements were made in benzene soln. for the mixed solutes: (1) Toluene-ethylbenzene: M is independent of the concn. of ($A + B$), for the pure components as well as for the mixt.; there is only one single isocoon. line, a straight line. (2) Methanol-ethanol: for all ratios $A:B$, the M increases with increasing concn. of ($A + B$); the isocoon. lines are nearly parallel straight lines. (3) Piperidine-phenyl isothiocyanate: all isocoon. lines show a common sharp max. at $A:B = 1$, which indicates compd. formation (phenylpiperidine-thiourea, isolated and identified). (4) Methylamine-phenyl isothiocyanate: same shape as foregoing, max. at $A:B = 1$. Systems 1 and 2 illustrate cases without chem. interaction between A and B . Moreover, in system 1, A and B are free from assocn. In system 2, both A and B are partly assoc.; the behavior of the M curves indicates that either the degree of assocn. of A and of B must remain unchanged with changing concn. or that disocn. of A_1 and of B_1 is compensated by an assocn. AB . Systems 3 and 4 are examples of components with chem. interaction. The compd. AB is assoc., the degree of assocn. rising with increasing concn. N. T.

N. T.

Mol. wt. determinations as a means of physico-chemical analysis.
V. V. Udovenko (*Acta Physicochim. U.R.S.S.*, 1942, 18, 236-347).---
The variation of mol. wt. with composition is determined by the
cryoscopic method and plotted in the case of MeOH-EtOH, and
PhMe-PhEt mixtures in C_6H_6 , to represent mixtures of similar
mols. where there is no chemical interaction, and for piperidine-
PhNCS and NtMePh-PhNCS mixtures, where there is interaction.
The mol. wt.-composition curve is a straight line where there is
no interaction, but the curve shows a max. coinciding with the com-
position of the compound where there is interaction. A. J. M.

2

The maximum of viscosity of irrational systems. V. V. Uglavanko (Phys. chem. Lab., Middle Asiatic State Univ.). Akad. Nauk S.S.S.R., Otdel. Tekh. Nauk, Inst. Mashinostroyeniya, Sovetskoye Vuzochiye Zhidkostey i Mashinostroyeniya (Conf. on Viscosity of Liquids and Colloidal Solns.) 2, 89-93 (1944).—Comparison with melting diagrams has shown that in "irrational" binary systems the max. of the viscosity curves in terms of compn. does not occur at the compn. corresponding to that of the compl. formed. The usual interpretation in terms of disson. of the compl. is subject to caution. A diagram of a system A-B involving a compl. C can be constructed from the part diagram A-C, with a max., and constructed from the part diagram B-C, in the resulting A-B C-B. The displacement of the compl. C and the concurrence is due to interaction of the compl. C and the component A and not to disson. This is further borne out by the observation that the max. does not tend to draw closer to the stoichiometric compn. as the temp. approaches the m.p., which would have to be the case if the displacement were due to disson. in liquid phase. The shift sometimes is found to persist below the m.p. in undercooled liquid. In the system $\text{SnCl}_4\text{-CH}_3\text{COOH}$, the max. cooled liquid. In the system $\text{SnCl}_4\text{-CH}_3\text{COOH}$, the max. of viscosity lies at about 25% SnCl_4 when the compl. has of viscosity 1:2 (m.p. 18.7°). In $\text{HClO}_4\text{-CH}_3\text{COOH}$, the compl. 1:2 (m.p. 67 mod. % CH_3COOH , and its position in the max. lies at 67 mod. % CH_3COOH , and its position is unaffected by temp., whereas the compl. (m.p. 41°) is equimol. The same is true for the system phenol-aniline; the compl. (m.p. 30.5°) is equimol., but the max. of viscosity is at 66.67 mod. % phenol, independently of temp. A further example of a shifted max. is $\text{HClO}_4\text{-H}_2\text{O}$. Compn. and stability of binary compds. were further investigated by cryoscopic measurements on benzene solns. of various concns. of both the pure components and their mixts. and the results represented graphically. The curves of cryoscopic mol. wt. for the system chloral-hydroxyl show the compl. to be stable at all dilns. and disson. to be absent. The max. on the curve corresponding to infinite diln. is singular (discontinuous); on the other hand, with increasing concn. of the benzene soln., the max. of mol. wt. shifts gradually towards the EIOH; this evidences interaction between that component and the ethereal ethylate. With rising temp., the max. moves towards the compl. In the 2 systems phenol-aniline and phenol-pyridine, the compl. is dissolved at high dilns.; at a concn. of about 1 mole/1000 g. benzene, disson. is practically zero. At higher concns., the cryoscopic mol. wt. shows a max., shifted towards phenol; this again indicates interaction between the phenol and the equimol. compl. Some cases are also known where the compl. exists in the liquid phase only and therefore is not expressed in the melting diagram but only in the viscosity curve (example the system $(\text{C}_6\text{H}_5)_3\text{CH-SbBr}_3$).

N. Thon

ABR-15A METALLURGICAL INFORMATION CLASSIFICATION
FROM SYMBOLISM

ISSUED WITH ONLY ONE

CLASSIFIED ON GHS 151

157 AND 158 CROCKS

159 AND 160 CROCKS

161 AND 162 CROCKS

163 AND 164 CROCKS

165 AND 166 CROCKS

167 AND 168 CROCKS

169 AND 170 CROCKS

171 AND 172 CROCKS

173 AND 174 CROCKS

175 AND 176 CROCKS

177 AND 178 CROCKS

179 AND 180 CROCKS

181 AND 182 CROCKS

183 AND 184 CROCKS

185 AND 186 CROCKS

187 AND 188 CROCKS

189 AND 190 CROCKS

191 AND 192 CROCKS

193 AND 194 CROCKS

195 AND 196 CROCKS

197 AND 198 CROCKS

199 AND 200 CROCKS

201 AND 202 CROCKS

203 AND 204 CROCKS

205 AND 206 CROCKS

207 AND 208 CROCKS

209 AND 210 CROCKS

211 AND 212 CROCKS

213 AND 214 CROCKS

215 AND 216 CROCKS

217 AND 218 CROCKS

219 AND 220 CROCKS

221 AND 222 CROCKS

223 AND 224 CROCKS

225 AND 226 CROCKS

227 AND 228 CROCKS

229 AND 230 CROCKS

231 AND 232 CROCKS

233 AND 234 CROCKS

235 AND 236 CROCKS

237 AND 238 CROCKS

239 AND 240 CROCKS

241 AND 242 CROCKS

243 AND 244 CROCKS

245 AND 246 CROCKS

247 AND 248 CROCKS

249 AND 250 CROCKS

251 AND 252 CROCKS

253 AND 254 CROCKS

255 AND 256 CROCKS

257 AND 258 CROCKS

259 AND 260 CROCKS

261 AND 262 CROCKS

263 AND 264 CROCKS

265 AND 266 CROCKS

267 AND 268 CROCKS

269 AND 270 CROCKS

271 AND 272 CROCKS

273 AND 274 CROCKS

275 AND 276 CROCKS

277 AND 278 CROCKS

279 AND 280 CROCKS

281 AND 282 CROCKS

283 AND 284 CROCKS

285 AND 286 CROCKS

287 AND 288 CROCKS

289 AND 290 CROCKS

291 AND 292 CROCKS

293 AND 294 CROCKS

295 AND 296 CROCKS

297 AND 298 CROCKS

299 AND 300 CROCKS

301 AND 302 CROCKS

303 AND 304 CROCKS

305 AND 306 CROCKS

307 AND 308 CROCKS

309 AND 310 CROCKS

311 AND 312 CROCKS

313 AND 314 CROCKS

315 AND 316 CROCKS

317 AND 318 CROCKS

319 AND 320 CROCKS

321 AND 322 CROCKS

323 AND 324 CROCKS

325 AND 326 CROCKS

327 AND 328 CROCKS

329 AND 330 CROCKS

331 AND 332 CROCKS

333 AND 334 CROCKS

335 AND 336 CROCKS

337 AND 338 CROCKS

339 AND 340 CROCKS

341 AND 342 CROCKS

343 AND 344 CROCKS

345 AND 346 CROCKS

347 AND 348 CROCKS

349 AND 350 CROCKS

351 AND 352 CROCKS

353 AND 354 CROCKS

355 AND 356 CROCKS

357 AND 358 CROCKS

359 AND 360 CROCKS

361 AND 362 CROCKS

363 AND 364 CROCKS

365 AND 366 CROCKS

367 AND 368 CROCKS

369 AND 370 CROCKS

371 AND 372 CROCKS

373 AND 374 CROCKS

375 AND 376 CROCKS

377 AND 378 CROCKS

379 AND 380 CROCKS

381 AND 382 CROCKS

383 AND 384 CROCKS

385 AND 386 CROCKS

387 AND 388 CROCKS

389 AND 390 CROCKS

391 AND 392 CROCKS

393 AND 394 CROCKS

395 AND 396 CROCKS

397 AND 398 CROCKS

399 AND 400 CROCKS

401 AND 402 CROCKS

403 AND 404 CROCKS

405 AND 406 CROCKS

407 AND 408 CROCKS

409 AND 410 CROCKS

411 AND 412 CROCKS

413 AND 414 CROCKS

415 AND 416 CROCKS

417 AND 418 CROCKS

419 AND 420 CROCKS

421 AND 422 CROCKS

423 AND 424 CROCKS

425 AND 426 CROCKS

427 AND 428 CROCKS

429 AND 430 CROCKS

431 AND 432 CROCKS

433 AND 434 CROCKS

435 AND 436 CROCKS

437 AND 438 CROCKS

439 AND 440 CROCKS

441 AND 442 CROCKS

443 AND 444 CROCKS

445 AND 446 CROCKS

447 AND 448 CROCKS

449 AND 450 CROCKS

451 AND 452 CROCKS

453 AND 454 CROCKS

455 AND 456 CROCKS

457 AND 458 CROCKS

459 AND 460 CROCKS

461 AND 462 CROCKS

463 AND 464 CROCKS

465 AND 466 CROCKS

467 AND 468 CROCKS

469 AND 470 CROCKS

471 AND 472 CROCKS

473 AND 474 CROCKS

475 AND 476 CROCKS

477 AND 478 CROCKS

479 AND 480 CROCKS

481 AND 482 CROCKS

483 AND 484 CROCKS

485 AND 486 CROCKS

487 AND 488 CROCKS

489 AND 490 CROCKS

491 AND 492 CROCKS

493 AND 494 CROCKS

495 AND 496 CROCKS

497 AND 498 CROCKS

499 AND 500 CROCKS

501 AND 502 CROCKS

503 AND 504 CROCKS

505 AND 506 CROCKS

507 AND 508 CROCKS

509 AND 510 CROCKS

511 AND 512 CROCKS

513 AND 514 CROCKS

515 AND 516 CROCKS

517 AND 518 CROCKS

519 AND 520 CROCKS

521 AND 522 CROCKS

523 AND 524 CROCKS

525 AND 526 CROCKS

527 AND 528 CROCKS

529 AND 530 CROCKS

531 AND 532 CROCKS

533 AND 534 CROCKS

535 AND 536 CROCKS

537 AND 538 CROCKS

539 AND 540 CROCKS

541 AND 542 CROCKS

543 AND 544 CROCKS

545 AND 546 CROCKS

547 AND 548 CROCKS

549 AND 550 CROCKS

551 AND 552 CROCKS

553 AND 554 CROCKS

555 AND 556 CROCKS

557 AND 558 CROCKS

559 AND 560 CROCKS

561 AND 562 CROCKS

563 AND 564 CROCKS

565 AND 566 CROCKS

567 AND 568 CROCKS

569 AND 570 CROCKS

571 AND 572 CROCKS

573 AND 574 CROCKS

575 AND 576 CROCKS

577 AND 578 CROCKS

579 AND 580 CROCKS

581 AND 582 CROCKS

583 AND 584 CROCKS

585 AND 586 CROCKS

587 AND 588 CROCKS

589 AND 590 CROCKS

591 AND 592 CROCKS

593 AND 594 CROCKS

595 AND 596 CROCKS

597 AND 598 CROCKS

599 AND 600 CROCKS

601 AND 602 CROCKS

603 AND 604 CROCKS

605 AND 606 CROCKS

607 AND 608 CROCKS

609 AND 610 CROCKS

611 AND 612 CROCKS

613 AND 614 CROCKS

615 AND 616 CROCKS

617 AND 618 CROCKS

619 AND 620 CROCKS

621 AND 622 CROCKS

623 AND 624 CROCKS

625 AND 626 CROCKS

627 AND 628 CROCKS

629 AND 630 CROCKS

631 AND 632 CROCKS

633 AND 634 CROCKS

635 AND 636 CROCKS

637 AND 638 CROCKS

639 AND 640 CROCKS

641 AND 642 CROCKS

643 AND 644 CROCKS

645 AND 646 CROCKS

647 AND 648 CROCKS

649 AND 650 CROCKS

651 AND 652 CROCKS

653 AND 654 CROCKS

655 AND 656 CROCKS

657 AND 658 CROCKS

659 AND 660 CROCKS

661 AND 662 CROCKS

663 AND 664 CROCKS

665 AND 666 CROCKS

667 AND 668 CROCKS

669 AND 670 CROCKS

671 AND 672 CROCKS

673 AND 674 CROCKS

675 AND 676 CROCKS

677 AND 678 CROCKS

679 AND 680 CROCKS

681 AND 682 CROCKS

683 AND 684 CROCKS

685 AND 686 CROCKS

687 AND 688 CROCKS

689 AND 690 CROCKS

691 AND 692 CROCKS

693 AND 694 CROCKS

695 AND 696 CROCKS

697 AND 698 CROCKS

699 AND 700 CROCKS

701 AND 702 CROCKS

703 AND 704 CROCKS

705 AND 706 CROCKS

707 AND 708 CROCKS

709 AND 710 CROCKS

711 AND 712 CROCKS

713 AND 714 CROCKS

715 AND 716 CROCKS

717 AND 718 CROCKS

719 AND 720 CROCKS

721 AND 722 CROCKS

723 AND 724 CROCKS

725 AND 726 CROCKS

727 AND 728 CROCKS

729 AND 730 CROCKS

731 AND 732 CROCKS

733 AND 734 CROCKS

735 AND 736 CROCKS

737 AND 738 CROCKS

739 AND 740 CROCKS

741 AND 742 CROCKS

743 AND 744 CROCKS

745 AND 746 CROCKS

747 AND 748 CROCKS

749 AND 750 CROCKS

751 AND 752 CROCKS

753 AND 754 CROCKS

755 AND 756 CROCKS

757 AND 758 CROCKS

759 AND 760 CROCKS

761 AND 762 CROCKS

763 AND 764 CROCKS

765 AND 766 CROCKS

767 AND 768 CROCKS

769 AND 770 CROCKS

771 AND 772 CROCKS

773 AND 774 CROCKS

775 AND 776 CROCKS

777 AND 778 CROCKS

779 AND 780 CROCKS

781 AND 782 CROCKS

783 AND 784 CROCKS

785 AND 786 CROCKS

787 AND 788 CROCKS

789 AND 790 CROCKS

791 AND 792 CROCKS

793 AND 794 CROCKS

795 AND 796 CROCKS

797 AND 798 CROCKS

799 AND 800 CROCKS

801 AND 802 CROCKS

803 AND 804 CROCKS

805 AND 806 CROCKS

807 AND 808 CROCKS

809 AND 810 CROCKS

811 AND 812 CROCKS

813 AND 814 CROCKS

815 AND 816 CROCKS

817 AND 818 CROCKS

819 AND 820 CROCKS

821 AND 822 CROCKS

823 AND 824 CROCKS

825 AND 826 CROCKS

827 AND 828 CROCKS

829 AND 830 CROCKS

831 AND 832 CROCKS

833 AND 834 CROCKS

835 AND 836 CROCKS

837 AND 838 CROCKS

839 AND 840 CROCKS

841 AND 842 CROCKS

843 AND 844 CROCKS

845 AND 846 CROCKS

847 AND 848 CROCKS

849 AND 850 CROCKS

851 AND 852 CROCKS

853 AND 854 CROCKS

855 AND 856 CROCKS

857 AND 858 CROCKS

859 AND 860 CROCKS

861 AND 862 CROCKS

863 AND 864 CROCKS

865 AND 866 CROCKS

867 AND 868 CROCKS

869 AND 870 CROCKS

871 AND 872 CROCKS

873 AND 874 CROCKS

875 AND 876 CROCKS

877 AND 878 CROCKS

879 AND 880 CROCKS

881 AND 882 CROCKS

883 AND 884 CROCKS

885 AND 886 CROCKS

887 AND 888 CROCKS

889 AND 890 CROCKS

891 AND 892 CROCKS

893 AND 894 CROCKS

895 AND 896 CROCKS

897 AND 898 CROCKS

899 AND 900 CROCKS

901 AND 902 CROCKS

903 AND 904 CROCKS

905 AND 906 CROCKS

907 AND 908 CROCKS

909 AND 910 CROCKS

911 AND 912 CROCKS

913 AND 914 CROCKS

915 AND 916 CROCKS

917 AND 918 CROCKS

919 AND 920 CROCKS

921 AND 922 CROCKS

923 AND 924 CROCKS

925 AND 926 CROCKS

927 AND 928 CROCKS

929 AND 930 CROCKS

931 AND 932 CROCKS

933 AND 934 CROCKS

935 AND 936 CROCKS

937 AND 938 CROCKS

939 AND 940 CROCKS

941 AND 942 CROCKS

943 AND 944 CROCKS

945 AND 946 CROCKS

947 AND 948 CROCKS

949 AND 950 CROCKS

951 AND 952 CROCKS

953 AND 954 CROCKS

955 AND 956 CROCKS

957 AND 958 CROCKS

959 AND 960 CROCKS

961 AND 962 CROCKS

963 AND 964 CROCKS

965 AND 966 CROCKS

967 AND 968 CROCKS

969 AND 970 CROCKS

971 AND 972 CROCKS

973 AND 974 CROCKS

975 AND 976 CROCKS

977 AND 978 CROCKS

979 AND 980 CROCKS

981 AND 982 CROCKS

983 AND 984 CROCKS

985 AND 986 CROCKS

987 AND 988 CROCKS

989 AND 990 CROCKS

991 AND 992 CROCKS

993 AND 994 CROCKS

995 AND 996 CROCKS

997 AND 998 CROCKS

999 AND 1000 CROCKS

1001 AND 1002 CROCKS

1003 AND 1004 CROCKS

1005 AND 1006 CROCKS

1007 AND 1008 CROCKS

1009 AND 1010 CROCKS

1011 AND 1012 CROCKS

1013 AND 1014 CROCKS

1015 AND 1016 CROCKS

1017 AND 1018 CROCKS

1019 AND 1020 CROCKS

1021 AND 1022 CROCKS

1023 AND 1024 CROCKS

1025 AND 1026 CROCKS

1027 AND 1028 CROCKS

1029 AND 1030 CROCKS

1031 AND 1032 CROCKS

1033 AND 1034 CROCKS

1035 AND 1036 CROCKS

1037 AND 1038 CROCKS

1039 AND 1040 CROCKS

1041 AND 1042 CROCKS

1043 AND 1044 CROCKS

1045 AND 1046 CROCKS

1047 AND 1048 CROCKS

1049 AND 1050 CROCKS

1051 AND 1052 CROCKS

1053 AND 1054 CROCKS

1055 AND 1056 CROCKS

1057 AND 1058 CROCKS

1059 AND 1060 CROCKS

1061 AND 1062 CROCKS

1063 AND 1064 CROCKS

1065 AND 1066 CROCKS

1067 AND 1068 CROCKS

1069 AND 1070 CROCKS

1071 AND 1072 CROCKS

1073 AND 1074 CROCKS

1075 AND 1076 CROCKS

1077 AND 1078 CROCKS

1079 AND 1080 CROCKS

1081 AND 1082 CROCKS

1083 AND 1084 CROCKS

1085 AND 1086 CROCKS

1087 AND 1088 CROCKS

1089 AND 1090 CROCKS

1091 AND 1092 CROCKS

1093 AND 1094 CROCKS

1095 AND 1096 CROCKS

1097 AND 1098 CROCKS

1099 AND 1100 CROCKS

1101 AND 1102 CROCKS

1103 AND 1104 CROCKS

1105 AND 1106 CROCKS

1107 AND 1108 CROCKS

1109 AND 1110 CROCKS

1111 AND 1112 CROCKS

1113 AND 1114 CROCKS

1115 AND 1116 CROCKS

1117 AND 1118 CROCKS

1119 AND 1120 CROCKS

1121 AND 1122 CROCKS

1123 AND 1124 CROCKS

1125 AND 1126 CROCKS

1127 AND 1128 CROCKS

1129 AND 1130 CROCKS

1131 AND 1132 CROCKS

1133 AND 1134 CROCKS

1135 AND 1136 CROCKS

1137 AND 1138 CROCKS

1139 AND 1140 CROCKS

1141 AND 1142 CROCKS

1143 AND 1144 CROCKS

1145 AND 1146 CROCKS

1147 AND 1148 CROCKS

1149 AND 1150 CROCKS

1151 AND 1152 CROCKS

1153 AND 1154 CROCKS

1155 AND 1156 CROCKS

1157 AND 1158 CROCKS

1159 AND 1160 CROCKS

1161 AND 1162 CROCKS

1163 AND 1164 CROCKS

1165 AND 1166 CROCKS

1167 AND 1168 CROCKS

1169 AND 1170 CROCKS

1171 AND 1172 CROCKS

1173 AND 1174 CROCKS

1175 AND 1176 CROCKS

1177 AND 1178 CROCKS

1179 AND 1180 CROCKS

1181 AND 1182 CROCKS

1183 AND 1184 CROCKS

1185 AND 1186 CROCKS

1187 AND 1188 CROCKS

1189 AND 1190 CROCKS

1191 AND 1192 CROCKS

1193 AND 1194 CROCKS

1195 AND 1196 CROCKS

1197 AND 1198 CROCKS

1199 AND 1200 CROCKS

1201 AND 1202 CROCKS

1203 AND 1204 CROCKS

1205 AND 1206 CROCKS

1207 AND 1208 CROCKS

1209 AND 1210 CROCKS

1211 AND 1212 CROCKS

1213 AND 1214 CROCKS

1215 AND 1216 CROCKS

1217 AND 1218 CROCKS

1219 AND 1220 CROCKS

1221 AND 1222 CROCKS

1223 AND 1224 CROCKS

1225 AND 1226 CROCKS

1227 AND 1228 CROCKS

1229 AND 1230 CROCKS

1231 AND 1232 CROCKS

1233 AND 1234 CROCKS

1235 AND 1236 CROCKS

1237 AND 1238 CROCKS

1239 AND 1240 CROCKS

1241 AND 1242 CROCKS

1243 AND 1244 CROCKS

1245 AND 1246 CROCKS

1247 AND 1248 CROCKS

1249 AND 1250 CROCKS

1251 AND 1252 CROCKS

1253 AND 1254 CROCKS

1255 AND 1256 CROCKS

1257 AND 1258 CROCKS

1259 AND 1260 CROCKS

1261 AND 1262 CROCKS

1263 AND 1264 CROCKS

1265 AND 1266 CROCKS

1267 AND 1268 CROCKS

1269 AND 1270 CROCKS

1271 AND 1272 CROCKS

1273 AND 1274 CROCKS

1275 AND 1276 CROCKS

1277 AND 1278 CROCKS

1279 AND 1280 CROCKS

1281 AND 1282 CROCKS

1283 AND 1284 CROCKS

1285 AND 1286 CROCKS

1287 AND 1288 CROCKS

1289 AND 1290 CROCKS

1291 AND 1292 CROCKS

1293 AND 1294 CROCKS

1295 AND 1296 CROCKS

1297 AND 1298 CROCKS

1299 AND 1300 CROCKS

1301 AND 1302 CROCKS

1303 AND 1304 CROCKS

1305 AND 1306 CROCKS

1307 AND 1308 CROCKS

1309 AND 1310 CROCKS

1311 AND 1312 CROCKS

1313 AND 1314 CROCKS

1315 AND 1316 CROCKS

1317 AND 1318 CROCKS

1319 AND 1320 CROCKS

1321 AND 1322 CROCKS

1323 AND 1324 CROCKS

1325 AND 1326 CROCKS

1327 AND 1328 CROCKS

1329 AND 1330 CROCKS

1331 AND 1332 CROCKS

1333 AND 1334 CROCKS

1335 AND 1336 CROCKS

1337 AND 1338 CROCKS

1339 AND 1340 CROCKS

1341 AND 1342 CROCKS

1343 AND 1344 CROCKS

1345 AND 1346 CROCKS

1347 AND 1348 CROCKS

1349 AND 1350 CROCKS

1351 AND 1352 CROCKS

1353 AND 1354 CROCKS

1355 AND 1356 CROCKS

1357 AND 1358 CROCKS

1359 AND 1360 CROCKS

1361 AND 1362 CROCKS

1363 AND 1364 CROCKS

1365 AND 1366 CROCKS

1367 AND 1368 CROCKS

1369 AND 1370 CROCKS

1371 AND 1372 CROCKS

1373 AND 1374 CROCKS

1375 AND 1376 CROCKS

1377 AND 1378 CROCKS

1379 AND 1380 CROCKS

1381 AND 1382 CROCKS

1383 AND 1384 CROCKS

1385 AND 1386 CROCKS

1387 AND 1388 CROCKS

1389 AND 1390 CROCKS

1391 AND 1392 CROCKS

1393 AND 1394 CROCKS

1395 AND 1396 CROCKS

1397 AND 1398 CROCKS

1399 AND 1400 CROCKS

1401 AND 1402 CROCKS

1403 AND 1404 CROCKS

1405 AND 1406 CROCKS

1407 AND 1408 CROCKS

1409 AND 1410 CROCKS

1411 AND 1412 CROCKS

1413 AND 1414 CROCKS

1415 AND 1416 CROCKS

1417 AND 1418 CROCKS

1419 AND 1420 CROCKS

1421 AND 1422 CROCKS

1423 AND 1424 CROCKS

1425 AND 1426 CROCKS

1427 AND 1428 CROCKS

1429 AND 1430 CROCKS

1431 AND 1432 CROCKS

1433 AND 1434 CROCKS

1435 AND 1436 CROCKS

1437 AND 1438 CROCKS

1439 AND 1440 CROCKS

1441 AND 1442 CROCKS

1443 AND 1444 CROCKS

1445 AND 1446 CROCKS

1447 AND 1448 CROCKS

1449 AND 1450 CROCKS

1451 AND 1452 CROCKS

1453 AND 1454 CROCKS

1455 AND 1456 CROCKS

1457 AND 1458 CROCKS

1459 AND 1460 CROCKS

1461 AND 1462 CROCKS

1463 AND 1464 CROCKS

146

Investigations of systems with formic acid. I. V. V. Udevenko and R. P. Alrapatova. *J. Gen. Chem.* (U.S.S.R.) 17, 425-9 (1947) (in Russian).—Viscosity, d , and cond. of solns. of HCOOH (I) in pyridine (II), in nitrobenzene (III), and in AcOH (IV) were measured at 0°, 25°, and 80°. The viscosity-concn. curves of I-II solns. show max. at about 70 mole % I at all temps. The cond. exhibits a max. at about 80 mole % I at all temps. and an inflection point at about 75 mole % I at 80°, which becomes a slight min. when the temp. is lowered to 0°. The viscosity-concn. curve of I-III solns. at 0° is S-shaped, with a max. at 86 mole % I and a min. at 41 mole % I. The S-shape becomes much less pronounced as the temp. is raised. The cond.-concn. curves for these solns. are also S-shaped, with inflection points at about 65 mole % I and without max. or min. The only other system displaying this type of curve is the KCl-MgCl_2 system, in which two compds. have been found, viz., $\text{KCl} \cdot \text{MgCl}_2$ and $2\text{KCl} \cdot \text{MgCl}_2$ (cf. *C.A.* 29, 22347). The viscosity-concn. curve of I-IV solns. shows a pronounced max. at 0° at about 90 mole % I, which becomes less pronounced as the temp. is raised. The cond. curves for this system show no max. or min. and are convex towards the concn. axis. The types of chem. interaction responsible for these results are discussed. Arild J. Müller

COMMON ELEMENTS		COMMON VARIABLE MODS	
<p>Investigation of systems with formic acid. II. V. V. Udovenko and N. P. Almyetov. <i>J. Gen. Chem.</i> (U.S.S.R.) 17, 685-8 (1947) (in Russian); <i>cf. C.A.</i> 42, 2502a. -- Viscosity, elec. cond., and d. were studied in binary systems for formic acid (I) with Et formate (II) and with Et ether (III). No conopds. are formed in either system, but there is a weak interaction between I and II. In the I-II system the viscosity decreases smoothly from I to II, the curve being concave towards the concn. axis. Values of the viscosity at 0, 25, and 50° for I are 2.8210, 1.5372, and 0.9767; and for II, 0.5288, 0.2073, and 0.3080 centipoise. The cond.-concn. curves are similar, except that there are breaks in the curves at about 25 mole % II. Values of the cond. are: I, 0.74, 1.34, 1.76; and II, 0.003, 0.003, and 0.003×10^{-1} ohm⁻¹ cm.⁻¹. Values of the d. are: I 1.2275, 1.2068, 1.1846; and II 0.9474, 0.9168, 0.8818 g./cc. No breaks or max. were observed in the curves of d., viscosity, or cond. of I-III systems. The viscosity-concn. curve was similar to the I-II system. At 0 and 25°, III has values of: viscosity 0.2668 and 0.2461 centipoise; cond. less than 10^{-7} ohm⁻¹ cm.⁻¹ at both temps.; d., 0.7323 and 0.7048 g./cc. Arild J. Miller</p>		2	
<p>ASB-ELA REVALUATION LITERATURE CLASSIFICATION</p>			
<p>RIGHT SYMBOL</p>		<p>RIGHT SYMBOL</p>	
<p>RIGHT SYMBOL</p>		<p>RIGHT SYMBOL</p>	

UDOVENKO, V. V.

Determination of the molecular weight as a method of physicochemical analysis. II. V. V. Udoenko and S. V. Babak, *J. Gen. Chem.* (U.S.S.R.) 17, 656-64 (1947) (in Russian); cf. *C.A.* 35, 6175¹.—The apparent mol. wt. M of a few binary systems of noninteracting components one of which is subject to assocn. was detd. by cryoscopy in C_6H_6 soln. and was plotted against the compn. for the total molalities $m = 0$ (infinite diln.), 0.5, 1.0, and 1.5 moles/1000 g. C_6H_6 . In the system PhMe-MeOH , the $m = 0$ isoconcentrate is a straight line; the $m = 0.5$ curve is convex to the axis of compn., the $m = 1.0$ and 1.5 curves pass through a min. shifting to the PhMe side with rising m . In PhMe-EtOH , $m = 0$ is linear, the $m = 0.5$, 1.0, and 1.5 curves pass through a min. increasingly shifted to PhMe with rising m . In the PhMe-PhNO_2 , the $m = 0$ isoconcentrate is again rectilinear, the other curves convex to the axis of compn., merging with the $m = 0$ line at 72 mol. % PhMe . The system PhMe-PhNH_2 shows the same behavior; extrapolation to $m = 0$ gives for PhNH_2 a value of M somewhat higher than the theoretical; this was reproduced in repeated detns. irrespective of addnl. purification. The linearity of M at infinite diln. indicates absence of assocn. The min., and its shift, is the result, on the one hand, of the decrease of M due to decreasing assocn. of the alc. with its increasing diln. by PhMe , and, on the other hand, to the increase of M with increasing PhMe content. The absence of a min. in the case of PhNO_2 and of PhNH_2 means not only lower assocn. but mainly a lesser tendency to disassocn. on diln. with PhMe .

N. Thou

1ST AND 2ND PAGES PROCESS AND PROPERTIES INDEX 2

CA

Determination of the molecular weight as a method of physicochemical analysis. III. V. V. Udorenko and S. F. Babak (Mid-Asiatic State Univ., Tashkent), *J. Gen. Chem. (U.S.S.R.)* 17, 1563-6 (1947) (in Russian); *cf. C.A.* 42, 4384; 43, 6178. Apparent mol. wts. M (determ. by cryoscopy in C_2H_5 soln.) of a series of binary mixts. with $CHCl_3$ are tabulated and plotted as a function of the compn. of the mixt., for four concns.: infinite diln., 0.5, 1.0, and 1.5 moles/1000 g. C_2H_5 (isoconcentrate lines I-IV). (1) In $EtOH-CHCl_3$, I is rectilinear, II convex to the axis of abscissas, III and IV pass through a min.; the lines draw close with increasing $CHCl_3$ and merge with I at 93 mole % $CHCl_3$. The diagram indicates disson. of assocd. $EtOH$ moles, which is borne out by the neg. thermal effect in mixing $EtOH$ with a large amt. of $CHCl_3$. The chem. interaction indicated by the viscosity isotherms does not appear on the M diagram. (2) In $PhNH_2-CHCl_3$, lines I-IV have a shape similar to the corresponding lines of the preceding system; the lines merge with I at 68 mole % $CHCl_3$. The presence of the min. on III and IV, compared with its absence in the system aniline-toluene, indicates increased disson. of assocd. aniline moles in the presence of $CHCl_3$ as compared with toluene. (3) In $PhNO_2-CHCl_3$, all 4 lines are convex and merge at 80 mole % $CHCl_3$; this indicates only disson. (4) $Me_2CO-CHCl_3$: I is rectilinear, III-IV are slightly concave; this indicates some chem. interaction. N. Thon

ASS-55A METALLURGICAL LITERATURE CLASSIFICATION

1ST AND 2ND PAGES PROCESS AND PROPERTIES INDEX

UDOVENKO, V. V., PROF.

Solution (Chemistry)

Some problems of investigating liquid binary systems. Biul.Sredneaz.un.,
no. 25, 1947.

9. Monthly List of Russian Accessions, Library of Congress, November 1952, Unclassified.

UDOVENKO V. V.

PA 21/49T13

USSR/Chemistry - Heat of Vaporization Oct 48
Chemistry - Mixture

"On Heats of Vaporization of Binary Mixtures, III,"
V. V. Udoenko, Ts. B. Frid, Gen Akad State U,
Lab Phys Chem, Tashkent, 8 pp

"Zhur Fiz Khimii" No 10

Studies vapor pressure of five systems formed by
dichloroethane with alcohols: methyl and ethyl at
40, 50 and 60°, and propyl, isobutyl, and isooctyl
at 50, 60, 70 and 80°. Calculates heats of
vaporization of these systems at one temperature
by own formula. Results agree with calculations
using Clapeyron-Clausius equation. Shows that own
IC 21/49T13

USSR/Chemistry - Heat of Vaporization Oct 48
(Contd)

Formula can also be used to calculate heats of
vaporization of real binary liquid systems.
Submitted 6 Dec 47.

IC

21/49T13

UDEVENKO, V. V.

PA 8/49T29

USSR/Chemistry - Analysis, Physicochemical Apr 48
Chemistry - Molecular Weight, Determination of

"Determination of Molecular Weight as One Method of
Physicochemical Analysis, IV," V. V. Udevenko,
S. F. Babak, Lab Phys Chem, Gen Asiatic State U,
64 pp

"Zhur Obshoh Khim" Vol XVIII (LXXX), No 4

Studies following systems by molecular weight method:
methyl alcohol-aniline, methyl alcohol-nitrobenzene,
ethyl alcohol-aniline and ethyl alcohol-nitrobenzene.
Analyzes molecular weight diagrams of systems with
an associated component. Submitted 2 Dec 1946.
8/49T29

8/49130

USSR/Chemistry - Analysis, Physicochemical Apr 48
Chemistry - Molecular Weight, Determination
of

"Determination of the Molecular Weight as One Method
of Physicochemical Analysis, V," V. V. Udovenko,
B. F. Babak, Lab Phys Chem, Gen Asiatic State U,
5 1/2 pp

"Zhur Obshch Khim" Vol XVIII (LXXX), No 4

Studies following systems by the molecular weight
method: ethyl alcohol-acetone, aniline-acetone,
aniline-nitrobenzene and ethyl alcohol-acetic acid.
Shows how molecular weight diagram alters when
interaction of the components occurs at the same time
8/49130

USSR/Chemistry - Analysis, Physicochemical Apr 48
(Contd)

as dissociation of the associated molecules. Sub-
mitted 2 Dec 1946.

V. V.

UDOVENKO,

8/49130

UDOVENKO, V. V.

PA 8/49T21

USSR/Chemistry - Analysis, Physicochemical Apr 48
Chemistry - Molecular Weight, Determination
of

"Determination of Molecular Weight as One Method of
Physicochemical Analysis, VI," V. V. Udoenko, Lab
Phys Chem, Gen Asiatic State U, 71 pp

"Zhur Obshch Khim" Vol XVIII (LXXX), No 4

Studies following systems by the molecular weight
method: Chloral-methyl alcohol, chloral-ethyl alcohol,
chloral-isopropyl alcohol and chloral-n-butyl alcohol.
Analyzes molecular weight diagrams of systems with
sharply defined chemical reaction of the components.
Shows that irrational maximum is explained, not by
8/49T33

USSR/Chemistry - Analysis, Physicochemical Apr 48
(Contd)

dissociation of the chloraloholate, but by its
reaction with the alcohol. Submitted 21 Dec 1946.

8/49T21

UDOVENKO, V. V.

V. V. Udovenko and Tz. B. Fried, The heats of evaporation of binary mixtures. I. P. 1136.

An equation has been derived for the heat of evaporation of binary liquid mixtures. For the calculation of this value, one has to know the heats of evaporation and the vapor pressures of the pure components and also their activity coefficients in the mixtures. The vapor pressure of three ideal systems have been studied at 50°, 60°, 70° and 80°: propyl alcohol - isobutyl alcohol, propyl alcohol - isoamyl alcohol and isobutyl alcohol - isoamyl alcohol.

Lab. of Physical Chemistry of the
Central Asiatic State University,
Tashkent

September 24, 1947

SO: Journal of Physical Chemistry (USSR) 22, No. 9, 1948

UDOVENKO, V. D.

V. D. Udovenko and Tz. B. Fried, Heats of evaporation of binary mixtures. II. P. 1135.

This work includes the results of investigation of the vapor pressure and heat of evaporation of five systems composed of methyl, ethyl, propyl, isobutyl and isocamyl alcohols. The vapor pressure was studied by the dynamic method. The composition of the liquid and the vapor were determined refractometrically.

Central Asiatic State University,
Tashkent
December 1, 1947

SO: Journal of Physical Chemistry (USSR) 22, No. 9, 1948

1ST AND 2ND ORDERS										3RD AND 4TH ORDERS									
PROCESSES AND PROPERTIES INDEX																			
<p>Heats of vaporization of binary mixtures. III. V. V. Udovskii and Ya. B. Frid. <i>Zhur. Fiz. Khim.</i> (J. Phys. Chem.) 22, 1948-70 (1948); cf. C.A. 43, 4704. The compn. and pressure P of the azeotropic vapor of 8 binary systems containing dichloroethane (I) was detd. at 40, 50, and 60° for MeOH and EtOH and at 40, 60, 70, and 80° for PrOH, <i>iso</i>-BuOH, and <i>iso</i>-AmOH. The P of I was 180, 233.5, 280, 493, and 678.2 mm. Hg at 40, 50, 60, 70, and 80°, resp. At 50°, when the mole fraction x of the alc. in the liquid was 0.1, 0.4, 0.7, and 1.0, its mole fraction in the vapor and P were: for MeOH 0.478 and 404.2, 0.603 and 493.2, 0.657 and 501.2, and 1.0 and 422.0; for EtOH 0.530 and 315.7, 0.423 and 350.2, 0.501 and 376.2, and 1.0 and 333.0; for PrOH 0.106 and 246.0, 0.195 and 238.0, 0.317 and 198.2, and 1.0 and 87.0; for <i>iso</i>-BuOH 0.072 and 224.0, 0.136 and 216.0, 0.234 and 178.1, and 1.0 and 66.0; and for <i>iso</i>-AmOH 0.018 and 219.0, 0.064 and 182.2, 0.086 and 150.0, and 1.0 and 17.5. P has a max. in all systems except I + <i>iso</i>-AmOH; it is observed at greater x the higher the temp. The heat of vaporization of these systems calcd. according to the earlier equation, cf. <i>loc. cit.</i>, agrees with that calcd. from the equation of Clapeyron and Clausius. The former equation can be used also for nonideal systems in which the coeff. of activity varies with temp. J. J. B.</p>																			
ASM-ELA METALLURGICAL LITERATURE CLASSIFICATION										131000 001000 001000 001000									
131000 001000 001000 001000										131000 001000 001000 001000									

UDOVENKO, V.V., TOROPOV, A.P. and OSININA, M.Ye.

Udovenko, V.V. - "Conductometric titration of anabasine," Doklady Akad. nauk UzSSR, 1949, No. 1, p. 7-10 --- Summary in Uzbek

SO: U-3566, 15 March, 53, (Letopis 'Zhurnal 'nykh Statey, No. 14, 1949).

UDOVENKO, V. V.

Ayrapetova, R. P., Granitova, C. I., and Udovenk, V. V. - "Physical-chemical investigation of a formic acid-phenol system", Doklady Akad. nauk UzSSR, 1949, No. 2, p. 13-20, (Resume in Azerbaijani), - Bibliog: 14 iters.

SO: U-4392, 19 August 53, (Letopis 'Zhurnal 'nykh Statey, No. 21, 1949).

UDOVENKO, V. V.

"Research on a system with formic acid IV." Udovenko, V. V. and Airapetova, R. P.
(p.632)

SO: Journal of General Chemistry, (Zhurnal Obshchei Khimii) 1949, Vol. 19, No. 4.

UDOVENKO, V. V.

Chemical Abst.
Vol. 48 No. 4
Feb. 25, 1954
Inorganic Chemistry

Complex compounds of anabasine with cobalt salts.
T. I. Sushkevich and V. V. Udoenko (Mid-Asiatic State
Univ., Tashkent). *Doklady Akad. Nauk USSR*, S.S.S.R.
1949, No. 3, 18-20 (in Russian).—Addn. of anabasine to aq.
CoCl₂, with cooling, followed by addn. of HCl to dissolve the
greenish ppt. of hydrated Co, gave blue CoCl₂.C₁₀H₁₁N₃.
2HCl, m. 283°, when an excess of HCl was used, or the cor-
responding CoCl₂.C₁₀H₁₁N₃.HCl, m. 303°, with the corres-
pondingly smaller amount of HCl. Both are sol. in H₂O,
insol. in usual org. solvents, except for hot HCO₂H; both
are electrolytes in aq. soln., and the cond. indicates complete
dissoen. in soln. of the di-HCl salt with 6 ions being formed;
the mono-HCl salt is a weak electrolyte. Anabasine also
forms complexes with CuCl₂ and MnCl₂, which are not de-
scribed. G. M. Kosolapoff

MF
7-27-54

2

CA

Determination of molecular weight as a method of physical-chemical analysis. V. Y. Vukobratovich (Central Asia State Univ., Tashkent, U.S.S.R.). *Izv. Akad. Nauk Khim. Anal. Inst. Obshch. i Neorg. Khim. Akad. Nauk S.S.S.R.* 17, 101-10 (1946). The use of the method is illustrated on MeOH and EtOH in C_6H_6 . On the abscissa is plotted the concn. of the alc. in C_6H_6 expressed as mole. of alc. per 100 g. of C_6H_6 and on the ordinate the apparent mol. wt. A family of curves is obtained of which the lowest curve is that of MeOH and the upper one of EtOH at each concn. in C_6H_6 . Between these two are the curves of alc. mixes. taken in various proportions but of the same total concn. in C_6H_6 . Extrapolation to the ordinate, i.e. at zero concn. in C_6H_6 , gives the mol. wt. of infinitely dil. soln. On the basis of this diagram is plotted a diagram of compn.-mol. wt. on which MeOH-EtOH compn. is the abscissa and the mol. wt. the ordinate. The family of curves (straight lines) thus obtained are isoconcentrates in which the lowest is at zero concn. and the others at increasingly higher concns. By this method were obtained compn.-mol. wt. curves for a no. of systems. In the system toluene-ethylbenzene, the components do not react with each other and the mol. wt. of the pure components as well as of their mixt. does not change with their concn. in C_6H_6 . Therefore, all the isoconcentrates coincide and on the diagram appears one straight line. In the system toluene-MeOH, the latter is an associ. compd. and its degree of assocn. depends on its concn. in C_6H_6 or in its mixt. with toluene. Thus, at zero concn. of MeOH, the isoconcentrate is a straight line and as its quantity increases the curves bend toward the abscissa. Similarly are discussed the diagrams of toluene-aniline, nitrobenzene-EtOH, acetone-EtOH, phenol-pyridine, formic acid-EtOH, methylaniline-phenylisothiocyanate, and chloral-EtOH. M. Hosh

2

CA

Systems with formic acid. V. V. Usovskii and R. P. Alapetova. *Zhur. Obshch. Khim.* 19, 127-34; *J. Gen. Chem. U.S.S.R.* 19, 142-5(1946) (English translation); *Ch. C.A.* 42, 2680b.—Mol. wt. were detd. for various mixts. of HCOOH (I) with PbNO_3 (II), AcOH (III), HIO_3CH (IV), and Bi_2O_3 (V). The solvent used in the mol. wt. detns. was C_2H_5 . Graphs are given showing mol. wt. vs. concn. of I (in mole %) for 4 different total concns. of dissolved substances, viz., 0.0, 0.5, 1.0, and 1.5 moles per 1000 g. of C_2H_5 . The values graphed were extrapolated or interpolated from the exptl. results. In all cases the curves are displaced towards higher mol. wt. by increasing the total concn. In the system I-II, the mol. wt.-concn. curve for the 0.5 motal case is linear, extending from (approx.) 96 for I to 120 for II. In the other three cases the curves are linear in the range 0-80 mole % I. From 80 to 100 mole % I, the 0.0 motal curve is concave, and the 1.0 and 1.5 motal curves are convex, toward the concn. axis, giving values for the mol. wt. of I of, resp., 84, 95, and 115. In the system I-III, all the mol. wt.-concn. curves are linear, the 0.5 motal curve, e.g., extending from a mol. wt. of about 97 for I to about 128 for III. With increasing total concn. the slopes of the lines decrease. In the systems I-IV and I-V the curves all show max. In the I-IV system, the 0.5 motal curve rises from a mol. wt. of 96 for I to a max. of 100 at 80 mole % I, and then drops to 78 for IV. The sharpness of the max. increases with increasing total concn. In the I-V case, the 0.5 motal curve rises from a mol. wt. of 96 for I to a max. of 105 at 80 mole % I and then drops to 78.3 for V. The mol. wt. of pure V changes less with concn. than that of any other pure substance or soln. investigated, and there is no consistent trend, whereas the mol. wt. of all other substances increased with increasing total concn.

Artid J. Miller

UDOVENKO, V. V.

58/49124

USSR/Chemistry - Chloral
Chemistry - Viscometric Analysis Jan 49

"Viscosity of Systems of Chloral With Alcohols,"
V. V. Udoenko, Ye. I. Kalavenovskaya, M. P.
Prokopyeva, Lab of Physicochem, Cen Asiat'c
State U, 3 1/4 pp

"Zhur Obshch Kniz" Vol XIX, No 1

Tabular studies of viscosities and densities
of the system of chloral and methyl alcohol
at 40, 60, and 75° and the system of chloral
and isocetyl alcohol at 40, 60, and 80° show
maximum viscosity of the systems is invariably

58/49124

USSR/Chemistry - Chloral (Contd) Jan 49
affected by the alcohol, regardless of the
value for the viscosity of the chloral. Sub-
mitted 29 Sep 47.

58/49124

UDOVENKO, V. V.

PA 67/49T44

USSR/Chemistry - Anabesine
Acids, Mineral

May 49

"Reaction of Anabesine With Mineral Acids," V. V.
Udovenko, L. A. Vvedenskaya, Lab of Physicochem,
Cen Asia State U, 1 3/4 pp

"Zhur Obshch Khim" Vol XIX, No 5

Derives the crystalline compounds of anabesine with
hydrogen chloride and nitric acid, respectively:

$C_{10}H_{14}N_2 \cdot HCl$ and $C_{10}H_{14}N_2 \cdot HNO_3$. Submitted
28 Mar 48.

67/49T44

CA

Systems with formic acid. IV. V. V. Ushakov and
N. P. Alapetova. *Zhur. Obshch. Khim.* 19, 632-6;
J. Gen. Chem. U.S.S.R. 19, 663-6 (1946) (English transla-
tion); cf. preceding abstr.—Mol. wt. were detd. for
various mixts. of formic acid (I) with acetone (II), 2-
butanone (III), and 2-pentanone (IV). As in the pre-
vious article, curves are shown for mol. wt. vs. concn. for
total concn. of dissolved substances equal to 0.0, 0.5, 1.0,
and 1.5 moles per 1000 g. benzene. In the system I-III,
the 0.0 and 0.5 molal curves are concave towards the
concn. axis, whereas the 1.0 and 1.5 molal curves go
through a max. at 90 mole % I. For the 1.0 molal case,
e.g., values of concn. of I (mole %) and mol. wt. are,
resp.: 100, 107; 90, 108; 80, 102; 70, 98; 60, 93. In
the system I-III, the curves at all total concns. go through
a max. at about 75 mole % I, the 1.0 molal curve going
from a mol. wt. of 107 for I, through a max. of 110, then
dropping to 74 for III. The system I-IV is similar, the 1.0
molal curve rising from a mol. wt. of 107 for I to a max. of
115 at 70 mole % I, and dropping to 89 for IV. The max.
in these curves are attributed to chem. interaction among
the components in the various systems. A. J. M.

UDOVENKO, V.V

Viscosimetry in physicochemical analysis. Trudy SAGU no.15:3-16
'50. (MLRA 9:5)

(Viscosimetry) (Chemistry, Analytical)

2A

Viscosity and density of binary systems with nicotine. " III. S. P. Babak and V. V. Ulovenko (Samarkand Med. Inst.). *Zhur. Obshchei Khim.* (J. Gen. Chem.) 20, 1598-71 (1950); cf. *C.A.* 44, 6220i. The nicotine-*o*-nitrophenol system shows a compl., *nicotine-2C₆H₄(OH)NO₂*. Viscosity of the system shows a max. at the above compl.; the highest max. is at 35° 8.0113 centipoises, lower max. at 50° 4.3888, and at 75° 2.0000. Density-compl. curves are smooth curves of linear type going from 0.9006 at 35° and 100 mol.% nicotine to 1.2513 at 15.63 mol.% nicotine, at 50° from 0.9806 to 1.2851 at 0.0% nicotine, and at 75° from 0.9670 to 1.2571 at 0.0% nicotine. The nicotine-PhNO₂ system shows a linear viscosity relation: at 25° ranging from 1.7687 (0.0% nicotine) to 3.8042, at 50° from 1.1925 to 2.0376, at 75° from 0.8406 to 1.2626; values of d. similarly range from 1.3003 to 1.0068, from 1.1718 to 0.9806, and from 1.1480 to 0.9670, resp. The system with *o*-MeC₆H₄NO₂ has linear viscosity relations: at 25° from 2.0887 (0% nicotine) to 3.8042 (100%), at 50° from 1.3477 to 2.0376, at 75° from 0.9730 to 1.2626; d. varies similarly: from 1.1578 to 1.0068, from 1.1313 to 0.9806, from 1.1000 to 0.9670, resp. The system with *p*-MeC₆H₄NO₂ has a linear viscosity relation: at 25° from 2.8023 (41.68 mol.% nicotine) to 3.8042 (100%), at 50° from 1.2008 (0%) to 2.0376 (100%), at 75° from 0.8088 (0%) to 1.2626 (100%); d. varies similarly: from 1.0703 to 1.0068, from 1.1223 to 0.9806, from 1.1013 to 0.9670, resp. G. M. K.

CA

2

The viscosity and density of binary systems with nicotine. III. S. P. Babak and V. V. Ukharenko (Samarkand Med. Inst.). *J. Gen. Chem. U.S.S.R.* 20, 1029-32 (1970). (Engl. translation). See *C.A.* 45, 1000. R. M. S.

CA

Viscosity of binary solutions with nicotine. IV. S. F. Babak and V. V. Udoenko (Sumarkand Med. Inst.). *Zhur. Obshchei Khim.* 20: 2199-2201, 2203-5 (Engl. translation); cf. *C.A.* 46, 4972d. η and d were detd. for solns. of nicotine (I) with CCl_4 at temps. of 25, 35, and 50°, and with 1,2-dichloroethane, $PhCl$, and $PhBr$ at temps. of 25, 50, and 75°. There is no indication of any chem. interaction between the components in any case. The η -concn. curves in each of the 4 systems is a straight line at the highest temp. studied, but becomes increasingly convex towards the concn. axis with decreasing temp. In the system I- CCl_4 , e.g., values of mole % I, d , and η , resp., are: 25°: 0.00, 1.5782, 0.9240; 25.80, 1.5773, 1.5459; 50.36, 1.5226, 2.2747; 75.70, 1.4034, 3.0763; 100.00, 1.0069, 3.8943; at 50°: 0.00, 1.5201, 0.9667; 37.70, 1.2690, 1.3156; 75.70, 1.0773, 1.7311; 100.00, 0.9860.

2.0870. Similar values at 25° for the remaining systems are: I-1,2-dichloroethane: 0.00, 1.2846, 0.7645; 25.31, 1.1493, 1.2300; 50.00, 1.0636, 1.8280; 77.10, 1.0328, 2.8510; I- $PhCl$: 0.00, 1.0081, 0.7124; 25.87, 1.0678, 1.3057; 50.63, 1.0439, 1.8380; 72.10, 1.0237, 2.8972; I- $PhBr$: 0.00, 1.4930, 1.0619; 25.14, 1.3301, 1.5435; 48.72, 1.2074, 2.1000; 74.67, 1.0916, 2.9670. V. 194. 2124-6. η and d were detd. at 25, 35, and 50° for the system I-acetone, and at 25, 50, and 75° for the system I-MeCOEt and nicotine-acetophenone (II). The first 2 systems have η -concn. curves similar to those mentioned in part IV except that even at the highest temp. investigated the curves are still convex towards the concn. axis. For the system I-II the curves are S-shaped, with the curvature decreasing with increasing temp. Values for mole % nicotine, d , and η , resp., at 25° are as follows: I-acetone: 0.00, 0.7818, 0.3332; 24.96, 0.8873, 0.6975; 50.03, 0.9448, 1.2707; 74.85, 0.9797, 2.2737; 100.00, 1.0098, 3.8942; I-MeCOEt: 0.00, 0.8804, 0.4004; 24.63, 0.8922, 0.7167; 48.78, 0.8385, 1.2450; 73.74, 0.9861, 2.8902; I-II: 0.00, 1.0242, 1.6170; 24.26, 1.0222, 2.1076; 49.34, 1.0162, 2.9460; 74.31, 1.0118, 3.6271.

Arild J. Møller

PASOVSKAYA, G.B.; UDOVENKO, V.V.

Simplified methods for the determination of alkali metals in presence
of magnesium. Trudy SAGU no.27:81-88 '51. (MLBA 9:5)
(Alkali metals) (Conductionmetric analysis)

PASOVSKAYA, G.B.; UDOVENKO, V.V.

Conductometric simultaneous determination of magnesium and alkali
metals. Trudy Kom. anal. khim. 4:196-204 '52. (MIRA 11:6)

(Magnesium)

(Alkali metals)

(Conductometric analysis)

UDOVENKO, V. V.

Chemical Abst.
Vol. 48 No. 8
Apr. 25, 1954
Analytical Chemistry

(3) 4
Conductometric determination of potassium with sodium
picrate. V. V. Udoenko and G. B. Pasovskaya (Central
Asian State Univ., Tashkent). *J. Anal. Chem. (U.S.S.R.)*
7, 181-2 (1952) (Engl. translation).—*Sci. C.A.* 47, 1636t.
H. L. H.

MF
11-4-54

UDOVJENKO, V., FATULINA, L. G.

Solubility

Solubility in the system ethyl alcohol - 1,2-ethylene dichloride - water. Zhur.fiz.khim., 16, No. 6, 1952.

Monthly List of Russian Accessions, Library of Congress, November 1952. Unclassified.

UDOVENKO, V. V.

Udovenko, V. V., Airapetova, R. P., Malakhova, V. T.- "Physico-chemical analysis of the systems: phenol-monochloro-acetic acid, and phenol-trichloroacetic acid." (p. 1759)

SO: Journal of General Chemistry, (Zhurnal Obshchei Khimii), 1952, Vol. 22, No. 10

Chemical Abst.
Vol. 48 No. 9
May 10, 1954
General and Physical Chemistry

5
③ Chem
/ Physicochemical analysis of the systems phenol-mono-
chloroacetic acid and phenol-trichloroacetic acid. V. V.
Udovenko, R. P. Alapetova, and W. T. Malakhova. J.
Gen. Chem. (U.S.S.R.) 22, 1801-2 (1952) (Engl. translation).
—See C.A. 47, 2028f. H. I. H.

UDOVENKO, V.V.

Chem

Chem Abs V48
1-25-54
Inorganic Chemistry

Some compounds of pyridine with cupric chloride. V. V. Udoenko and E. M. Ojapova (Mol.-Aviat. State Univ., Tashkent). *Zhur. Obshch. Khim.* 27, 2035-7 (1952). Equimol. amts. of anhyd. CuCl_2 and $\text{C}_5\text{H}_5\text{N}$, in abs. alc. or dry Me_2CO form light-blue $\text{CuCl}_2 \cdot 2\text{C}_5\text{H}_5\text{N}$ (I), which slowly (in a few hrs.) changes into gray $\text{CuCl}_2 \cdot \text{C}_5\text{H}_5\text{N}$ (II). With HCl, I forms 1.2HCl , yellow prisms, and dark brown crystals of $3\text{CuCl}_2 \cdot 2\text{C}_5\text{H}_5\text{N} \cdot 2\text{HCl} \cdot 2\text{H}_2\text{O}$ (III). This reaction can take place in moist HCl vapor, but not in dry HCl gas. II is not sol. in org. solvents; in H_2O , it is decompd. into I and CuCl_2 . Heated above 200° , I loses $\text{C}_5\text{H}_5\text{N}$ becoming anhyd. CuCl_2 . III is obtained from aq. solns. confg. strictly stoichiometric amts. of CuCl_2 and $\text{C}_5\text{H}_5\text{N}$ and an excess of HCl; without excess HCl, only II is pptd., but it redissolves in excess HCl and, on evapn., gives III, which can be recrystd. from H_2O . In alc., III decomp. to I, and in Me_2CO to 1.2HCl , insol. in C_6H_6 , CCl_4 , etc. III m. $194-5^\circ$ (decompn.). In air, III loses H_2O and HCl, becoming $3\text{CuCl}_2 \cdot 2\text{C}_5\text{H}_5\text{N}$ (IV), a light-green powder, which cannot be obtained directly from CuCl_2 and $\text{C}_5\text{H}_5\text{N}$. In H_2O , IV decomp. to I. With HCl, it forms III. It is stable in air at room temp., decomp. above 190° , and m. $272-3^\circ$ (decompn.). Blue $\text{CuCl}_2 \cdot 5\text{C}_5\text{H}_5\text{N}$ was obtained with CuCl_2 and a large excess of $\text{C}_5\text{H}_5\text{N}$, within 10 days at a temp. not over $12-15^\circ$, faster at -10° . N. Thon

UDOVENKO, V. V.

Chemical Abst.
Vol. 48 No. 9
May 10, 1954
Inorganic Chemistry

Some compounds of pyridine with cupric chloride. V. V. Udovenko and E. M. Osipova. *J. Gen. Chem. U.S.S.R.* 22: 2142-50 (1952) (Engl. translation).--See C.A. 48: 4906. H. L. H.

9-2-54
JP

General Chemistry 12

CA

Solubility in the system ethyl alcohol-1,2-dichloroethane-water. V. V. Udovenko and L. G. Fatkulina (Sverdlovsk. Gosudarst. Univ., Tashkent). *Zhur. Fiz. Khim.* 26, 802-7 (1952).—Soly. of H_2O in $C_2H_5Cl_2$ was detd. between 10° (0.0837 wt.%) and 60° (0.516%). Soly. of $C_2H_5Cl_2$ in H_2O was, e.g., 0.877% at 23° and 1.389% at 72.5°. Over 60 ternary solys. were measured. E.g., EtOH 31.7%, H_2O 76.24% dissolved 1.480% $C_2H_5Cl_2$ at 25°; EtOH 31.5%, H_2O 47.7% dissolved 18.45% $C_2H_5Cl_2$ at 34°; EtOH 31.5%, $C_2H_5Cl_2$ 68.5% dissolved 12.14% H_2O at 43°; and EtOH 10.6%, $C_2H_5Cl_2$ 80.4% dissolved 1.07% H_2O at 44°. Also d. and μ of many solns. at 30° and 40° and d. of solns. at 60° and 60° were detd. The compn. of conjugated layers is listed. Thus, at 30° the liquid H_2O 92.7, EtOH 6.3, $C_2H_5Cl_2$ 1.0% was in equill. with H_2O 0.5, EtOH 1.0, $C_2H_5Cl_2$ 98.5% and 81.5, 39.5, 9.0% in equill. with 0.5, 2.2, 97.3%. At 40° 88.8, 9.9, 1.3% was in equill. with 0.5, 2.2, 97.3%. At 60° 80.8, 17.0, 2.2% was in equill. with 0.9, 4.2, 94.9% and 41.7, 39.1, 19.2% with 7.2, 22.0, 70.8%. At 60° 80.1, 7.9, 2.0% was in equill. with 0.5, 1.5, 97.7% and 41.0, 37.0, 12.0% with 7.7, 21.5, 70.8%. The calcd. compn. of the crit. solys. was at 30° H_2O 18.3, EtOH 34.3, $C_2H_5Cl_2$ 47.4% at 40° 19.2, 32.9, 47.9% and at 60° 19.0, 31.4, 49.0%. J. J. Biberman

UDOVENKO, V. V., FATKULINA, L. G.

Chemical Apparatus.

Apparatus for determination of pressure and composition of saturated vapor of layer-forming liquid systems. Zhur.fiz.khim. 26, no. 2, 1952.

9. Monthly List of Russian Accessions, Library of Congress, September 1952, Unclassified.

UDOVENKO, V. V.; FATKULINA, L. G.

Phase Rule and Equilibrium.

Vapor pressure of ternary systems. Part 1. System ethyl alcohol - 1,
2-dichloroethanebenzene. Zhur. fiz. khim. 26 no. 5, 1952.

9. Monthly List of Russian Accessions, Library of Congress, November 1952/2 Unclassified.

FA 242T3

USSR/Chemistry - Ternary Systems

Nov 52

"The Equilibrium of Liquid-Liquid in a Ternary System," V. V. Udovenko and I. G. Fatkulina, Central Asian State U, Tashkent

"Zhur Fiz Khim" Vol 26, No 11, pp 1569-1572

The authors examined the eq of L. A. Rotinyan and showed that in the case of a paraboloidal curve of sepn in ternary systems, the nodes can be parallel to the side of the triangle when the liquid phases are in equil. On the basis of the above-mentioned eq, the authors concluded that the critical point of mutual soly in ternary systems can also be located at the vertex of the binodal curve of sepn.

242T3

UDOVENKO, V.V.; PASOVSKAYA, G.B.

Conductance method for the study of the adsorption of electrolytes.
Trudy SAGU no.33:35-37 '52. (MLRA 9:5)
(Electrolytes) (Adsorption)

UDOVENKO, V.V.

USSR:

Reaction of lupinine and diethylaminoethanol with copper chlorides. V. V. Udoenko and L. R. Altkandova (Central Asia State Univ., Tashkent). *Sbornik State Obshch. Khim.* 2, 1120-3 (1953).—Lupinine and dry CuCl_2 in abs. EtOH form brown $\text{CuL}_{10}\text{NOH} \cdot \text{CuCl}_2$, decomp. 125° , which slowly decomposes on exposure to air. If twice the proportion of lupinine is used, there is obtained a green alcoholate $\text{CuL}_{10}\text{NOCuCl}$ (I), m. 215° (from $\text{CH}_2\text{CH}_2\text{Cl}$), while the evapd. filtrate yields lupinine HCl salt, m. 211.5° . I also forms on mixing abs. EtOH soln. of lupinine with CuCl_2 in contact with atm. O. Addn. of abs. EtOH soln. of $\text{Et}_3\text{N} \cdot \text{CH}_2\text{CH}_2\text{OH}$ to EtOH soln. of CuCl_2 gave green $\text{Et}_3\text{NCH}_2\text{CH}_2\text{O} \cdot \text{CuCl}_2$, m. 169° (from $\text{ClCH}_2\text{CH}_2\text{Cl}$), while evapn. of the soln. gave the amine HCl salt. Thus the original adduct is transformed into the alcoholate under action of excess amino alc., which results in loss of HCl and formation of I and amine HCl salt. G. M. Kozlovskii

UDOVENKO, V. V.

USSR

✓ Separation of mixture of alkaloids anabasine and lupinine.
V. V. Udovertko, O. I. Granulova, and L. A. Vvedenskaya
~~Central Asian State Univ., Tashkent. Sbornik State~~
~~Sbornik Khim. 2, 1124-6 (1953).~~—The sepn. is performed
through the Cu salts (cf. following abstr.) as follows. The
mixt. of known compn. of anabasine and lupinine is treated
with a highly concd. soln. of CuCl_2 and HCl or $\text{Cu}(\text{NO}_3)_2$ and
 HNO_3 , made up on the basis of theoretical calcn. to form
either $\text{CuCl}_2 \cdot \text{A} \cdot 2\text{HCl}$ or $\text{Cu}(\text{NO}_3)_2 \cdot 2\text{A} \cdot 4\text{HNO}_3$ (A = ana-
basine), with cooling. The order of addn. is immaterial.
Since the mixt. now contains a ppt. of Cu oxide, addnl. acid
is added to dissolve this (avoid excess of HNO_3). Traces of
kerosine from com. alkaloids are removed at this point by
boiling. The mixt. then yields cryst. salt of anabasine,
which treated with concd. NaOH gives the pure base, which
is purified by distn. *in vacuo*. The evapd. filtrate from the
salt is treated with concd. NaOH and extd. with C_6H_6 .
Evapn. of the ext. yields cryst. lupinine. The HCl salt
gives somewhat better results. Pure anabasine b. $107-10^\circ$,
 n_D^{20} 1.5420, d_4^{20} 1.0400; lupinine m. 69° . G. M. K.

UDOVENKO, V. V.

USSR.

✓ Reaction of anabasina with salts of copper. V. V. Udo-
venko and O. I. Granitova (Central Asia Sci. Res. Inst.,
Tashkent). *Sbornik Statei Obshchei Khim.* 2, 1127-9
(1953).—Anabasina (A) forms 2 salts with CuCl_2 and HCl
and with CuBr_2 and HBr , resp. With control of the re-
actant proportions there were obtained: yellow $\text{CuCl}_2 \cdot A \cdot$
 2HCl , decomp. 205° (in aq. soln. this dis-
solves into components on basis of cond. detas.); $\text{CuCl}_2 \cdot A \cdot \text{HCl}$, green,
decomp. $191-6^\circ$ (completely dissolved in aq. soln.); black
 $\text{CuBr}_2 \cdot A \cdot 2\text{HBr}$, decomp. 214° ; red-brown $\text{CuBr}_2 \cdot A \cdot \text{HBr}$,

decomp. 157° . Similarly was obtained blue-violet Cu
 $(\text{NO}_3)_2 \cdot A \cdot \text{HNO}_3$, decomp. 107° , and blue $\text{CuSO}_4 \cdot A \cdot$
 $\text{H}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$, which loses all H_2O at 100° . All dis-
solves in aq. solns.
G. M. Kozolapoff

AP
1107

UDOVENKO, V.V.

USSR

Interaction of anabasine with mercuric and mercurous chlorides. V. V. UDOVENKO, M. A. AZIZOV, and E. I. ALIMOV. Doklady Akad. Nauk Uzbe. S.S.R. 1954, No. 3, 38-41; Referat. Zhur., Khim. 1954, No. 27-217. HgCl₂ reacted with anabasine (I) in aq. soln. to form a white amorphous compd. HgCl₂·C₁₀H₁₂N₂ (II), at 211° (decompn.). The reaction proceeds with appreciable evolution of heat and unless cooled a colored compound is obtained. II was obtained by adding dropwise an aq. soln. of I to an aq. soln. HgCl₂ cooled in ice water. Upon heating to 110° II turned yellow. It is nonhygroscopic, dissolves poorly in water (slightly better in hot water), acetone, and EtOH, insol. in benzene and toluene. After boiling, aq. solns. of II have an alk. reaction. II dissolves in I, liberating metallic Hg; addn. of Me₂CO hastens this reaction. In water acidified with AcOH or strong mineral acids II dissolved forming a compd. of different compn. Specifically aq. HCl formed HgCl₂·C₁₀H₁₂N₂·2HCl, acicular crystals. Hg₂Cl₂ (III) reacted with I according to: Hg₂Cl₂ + C₁₀H₁₂N₂ + HgCl₂ → C₁₀H₁₂N₂. A comparative study was made of the reaction of I, pyridine (IV), and piperidine (V) with III. The reactions were carried out with an excess of base in the absence of moisture. Immediately following the mixing, after 10 min., and after 7 days, the extent of interaction of I with III was 7, 9, and 18.9%, resp. For IV and III it was 30, 75.66, and 81.27%, and for V and III it was 14, 14 and 16.96%, resp. In the last case, no detn. was made after 7 days. Thus, in its action on dry III, I is closer to V than to IV. In the presence of water, the interaction of I or IV with III does not exceed 1%.

5

U.S.

U. J. Venko, U. V.

5
18

Interaction of anabasine with aluminum chloride and

Reaction of anabasine with organic acids. V. V. Vlas-
venko and L. A. Vyedenskaya (Central Asia State Univ.,
Tashkent). *Zhur. Obshch. Khim.* 23, 1031-4 (1953).
Anabasine (I) and HCO_2H yield crystalline $\text{I} \cdot \text{HCO}_2\text{H}$, very
hygroscopic and sol. in org. solvents. I and AcOH gave
 $\text{I} \cdot \text{AcOH}$, m. 88° . PrCO_2H similarly gave $\text{I} \cdot \text{PrCO}_2\text{H}$, very
hygroscopic solid. I and $\text{iso-BuCO}_2\text{H}$ gave $\text{I} \cdot \text{iso-BuCO}_2\text{H}$,
very hygroscopic solid. With $(\text{CO}_2\text{H})_2$ there are formed:
 $2\text{I} \cdot \text{C}_2\text{O}_4\text{H}_2$, m. 210° , $\text{I} \cdot \text{C}_3\text{O}_4\text{H}_4$, m. 199° , and $\text{I} \cdot 2\text{C}_4\text{O}_4\text{H}_6$, m.
m. 81° ; all are very sol. in the usual solvents. Malonic acid
gave $2\text{I} \cdot \text{C}_3\text{O}_4\text{H}_4$, m. 86° . Succinic acid gave very hygro-
scopic $2\text{I} \cdot \text{C}_4\text{O}_4\text{H}_6$. Tartaric acid gave solid $2\text{I} \cdot \text{C}_4\text{H}_4\text{O}_6$.
Citric acid gave solid $2\text{I} \cdot \text{C}_6\text{H}_8\text{O}_7$. With naphthalene-2-
sulfonic acid there are formed: $\text{I} \cdot \text{C}_{10}\text{H}_7\text{SO}_3\text{H}$, m. 139° and
 $\text{I} \cdot 2\text{C}_{10}\text{H}_7\text{SO}_3\text{H}$, m. 121° .
G. M. Kosolapoff

MS gw

UDOVENKO, V.V.

Udovenko, V V

The strength of some leg in water and aragon

USSR

Heats of vaporization of three-component liquid mixtures.
V. V. Udalovskiy and L. O. Potkullup (Central Asia State
Univ., Tashkent). *Zhur. Fiz. Khim.* 27, 1063-7 (1953);
cf. *Chim.* 47, 6743i. The heats of vaporization (L) of
ternary mixts. contg. 10-30 mole % EtOH (I) with varying
proportions of benzene (II) and 1,2-dichloroethane (III)
were calcd. by means of the Clausius-Clapeyron equation
from exptl. data published earlier (*loc. cit.*). Values of L so
calcd. agree within 1% or less on the av. with those calcd.
by means of the equation $L = L_1 y_1 + L_2 y_2 + L_3 y_3$, where
 L_i is the heat of vaporization of the i th component and y_i
is its mole fraction in the gas phase. Both sets of values are
tabulated. The equation $L = (L_1 p_{10} x_1 + L_2 p_{20} x_2 + L_3 p_{30} x_3) / (p_{10} x_1 + p_{20} x_2 + p_{30} x_3)$, where p_i , x_i , and x_{i0} are
the vapor pressure in the pure state, the activity coeff.,
and the mole fraction of the i th component in the liquid
phase, resp., was derived.
J. W. Loweberg, Jr.

① *gaw*

LIDOVENKO, V V

USSR.

...iteration of a ... anabazine and

LIPOVCHENKO, I. V.

U S S R

Viscosity of some binary systems with nicotine. V. V. Chelobanov and S. K. Bakak. Trudy Steklovsk. Kavalersk. Univ. Khim. Ser. 5, No. 40, p. 161 (1954). Refrat. 2,247. Khim. 1954, No. 2418; cf. C.I. 45, 46d, 46, 75001g. The viscosity and d_4^{20} of mixts. of nicotine with CHCl_3 , $\text{C}_2\text{H}_5\text{Br}$, tetrachloroethane, EtOH , i - BuOH , AmOH , and cyclohexyl alc. were studied at 25, 30, and 75°. and of nicotine and MeOH at 25 and 50°. The nature of the relation between the viscosity values and molar compn. indicates interaction between nicotine and molar compn. indicators: bromide, tetrachloroethane, and Me , Et , and i - Bu alcs. The interaction of these substances with nicotine indicates their acidic nature in relation to nicotine as a base. The viscosity isotherm of nicotine mixt. with cyclohexyl alc. shows no reaction between the two. The acidic properties of the halogen derivs. increase with the no. of halogen atoms in the mol. and increases also upon the replacement of Cl by Br . In the case of alcs., an increase in the length of the hydrocarbon chain attenuates their interaction with nicotine.

M. Rosch

MA

UDOVENKO, V.V.

✓1842. Conductometric titration of certain alkaloids.
V. V. Udovenko and L. A. Vvedenskaya. *Izv.
Khim. ZN.* 1974, 20 (8), 684-689.—By conducti-
metric titration with naphthalene-2-sulphonic acid
in acetone solution, the alkaloids present in the
following mixtures were separately determined:
anabasine - lupinine, anabasine - nicotine, ana-
basine - nicotine - lupinine, also anabasine - lupinine
in the presence of aphylline and aphyllidine. The
method depends on the effect of acetone in increas-
ing the difference between the ionisation constants
of the bases. It can be applied to the determination
of alkaloids in technical products.

E. W. Rikhsaidi

(2)

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

Vdovanko - V.V.

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

UDOVENKO, V. V.

U.S. A. T. 1980
U.S. A. T. 1980
U.S. A. T. 1980

AUTHORS: Udovenko, V.V. and Fialkov, Yu. Ya. bcl

TITLE: Viscosity of Germanium Tetrachloride - Ethers Systems. (Vyazkost' Sistem Chetyrekhkhloristyy Germaniy - Efiry).

PERIODICAL: "Zhurnal Neorganicheskoy Khimii" (Journal of Inorganic Chemistry Vol.11, No.2, pp.434-438 (U.S.S.R.)). 1976

ABSTRACT: A study of the complex-forming ability of germanium tetrachloride would be useful in enabling a comparison to be made of the properties of various addition products of IV group element tetrahalides; the rules governing the formation of these products could also be deduced. In connection with this studies have been made of the viscosities of binary systems of which one component was always germanium tetrachloride, the others being various organic compounds. All materials were carefully dried to avoid hydrolysis of the tetrachloride.

Measurements were made of the viscosity and also the density of the following systems: germanium tetrachloride - acetoacetic - ester, germanium tetrachloride - anisole at 20, 30 and 40°C; germanium tetrachloride - dioxane at 25 and 40°C, and germanium tetrachloride - diethyl ether and germanium - tetrachloride - dimethyl sulphide at 20°C. It was found that in contrast to titanium and tin tetrachlorides, germanium tetrachloride does not form addition products with the above type of compounds; this is explained as being connected with the increase in the

Card 1/2

581
Viscosity of Germanium Tetrachloride - Ethers Systems. (Cont.)
screening of the central atom in germanium tetrachloride.
There are thirty-one references, of which fifteen are Russian.
One Figure and Five Tables.
The work was carried out at Kiev Polytechnical Institute.
Received 17 September, 1956.

Card 2/2

UDOVENKO, U.U.

Viscosity of binary systems with chloral. I. V. V.
 Aldonko and R. I. P. Aldonko (M. I. Khim. i. Tekhn. 1964, 1, 107)
 of the system $\text{C}_6\text{H}_5\text{CHO}$ and H_2O at 20, 30, and 40°C, and
 of $\text{C}_6\text{H}_5\text{CHO}$ and $\text{C}_2\text{H}_5\text{OH}$ at 20 and 30°C were data. The
 systems show component interaction although a cryst
 adduct with cyclohexanol was the only definitely isolable
 product, in this case the viscosity maximum was structural
 max. at 90 mole % aldehyde. An equimolar mixt. of com-
 ponents forms a 1:1 adduct, in 94% (cf. Somerford and
 Cronk, C.A. 42, 2316). The system with PhCH_2OH failed
 to yield a cryst. adduct but gave a clear max. in the vis-
 cosity curve at 70 mole % alc. J. M. Kosolapoff

Udovenko, V. V.

Viscosity of binary systems from chloral
Udovenko and R. I. Klementov, *Dokl. Akad. Nauk SSSR*, 1971, 221, 1474.
Abstracted from *Chem. Abstr.* 67: 1474 (1971).
63068 Viscosity of the system CCl_3CHO -PhOH at 40°, 60°, and 80°, and CCl_3CHO -o-Cresol, CCl_3CHO -m-MeC₆H₄OH, and CCl_3CHO -p-MeC₆H₄OH at 25°, 50°, and 75°, was detd. There was indication that CCl_3CHO forms complexes with the 4 phenols that disso. into components when the temp. rises. The complex is apparently formed through H bonding and are not near associates. The system CCl_3CHO -PhOH yields a little $\text{CCl}_3\text{CH}(\text{OH})\text{Ph}$, m. 203°, owing to condensation under the influence of traces of HCl. Similarly a small amt. of 2,5-(10⁶ Me) with $\text{CH}(\text{OH})\text{CCl}_3$, m. 147°, was isolated. The system of PhOH- CCl_3CHO shows an irrational viscosity max. at 25 mole % CCl_3CHO ; this indicated a compd. $\text{CCl}_3\text{CHO} \cdot 3\text{PhOH}$; the system with o-cresol has an irrational max. that does not yield any rational component proportion; with m-cresol the system has an irrational max. at 33 mole % CCl_3CHO , as does that with p-cresol.
O. M. Kovalapal

Chem

MT

Udovenko, V.V.
USSR/Physical Chemistry - Thermodynamics, Thermochemistry, B-8
Equilibria, Physical-Chemical Analysis, Phase Transitions.

Abs Jour : Referat Zhur - Khimiya, No 1, 1958, 414

Author : V.V. Udovenko, Yu.Ya. Fialkov.

Inst :

Title : Viscosity of Systems Germanium Tetrachloride - Ethers and Esters.

Orig Pub : Zh. neorgan. khimii, 1957, 2, No 2, 434-438

Abstract : The viscosity and density of binary systems composed by germanium tetrachloride (I) with ethyl acetate (II), anisole (III), dioxane (IV), diethyl ether (V) and dimethylsulfide (VI) were measured. The systems I - II and I - III were studied at 20, 30 and 40°, the system I - IV was studied at 25 and 40°, and the systems I - V and I - VI were studied at 20°. All the operations of preparing the solutions and carrying out the measurements were done under airtight conditions. Viscosity was measured

Card 1/2

USSR/Physical Chemistry - Thermodynamics, Thermochemistry, B-8
Equilibria, Physical-Chemical Analysis, Phase Transitions.

Abs Jour : Ref Zhur - Khimiya, No 1, 1958, 414

in a closed viscosimeter for volatile liquids described earlier (Toropov A.P., Zh. prikl. khimii, 1939, 12, 1744). The viscosity isotherms of the studied systems are convex with reference to the composition axis, which indicates the absence of a chemical interaction between the components. The authors connect the absence of interaction with an increase of the screening of the central atom in I.

Card 2/2

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R001857820003-7"

446

AUTHORS: Udovenko, V. V., and Khomenko, R. I.

TITLE: Viscosity of Binary Systems with Chloral.4 (Vyazkost' binarnykh sistem s khloraem. IV.)

PERIODICAL: Zhurnal Obshchey Khimii, 1957, Vol. 27, No. 1, pp. 37-40 (U.S.S.R.)

ABSTRACT: Investigations were conducted to determine the viscosity and density of binary systems consisting of chloral with acetophenone (boiling point 80.0° at 12 mm); benzophenone (b. p. 161.0° at 11 mm); benzaldehyde (b. p. 177.2° at 731 mm) and salicylaldehyde (b. p. 193.0° at 727 mm). The viscosity/density measurements were carried out at temperatures of 25, 50 and 75° and the results obtained are given in tables (for each mixture separately). The isothermal viscosity/density curves are either slightly convex or rectilinear, depending upon the temperature. None of the systems investigated showed any definite reactions between the components which could lead to the formation of any specific chemical compound. Four tables. There are 5 references, of which 4 are Slavic.

Card 1/2

446

Viscosity of Binary Systems with Chloral.4

ASSOCIATION: Central Asiatic State University and the Kiev Polytechnicum
(Sredneaziatskiy Gosudarstvennyy Universitet i Kiyevskiy
Politekhnicheskiy Institut).

PRESENTED BY:

SUBMITTED: January 3, 1956

AVAILABLE:

Card 2/2

UDOVENKO, V. V.

79-2-9/58

AUTHORS:

Udovenko, V. V., and Khomenko, R. I.

TITLE:

Viscosity of Binary Systems with Chloral. Part 5. (Vyazkost' binarnykh sistem s khloralem. V.)

PERIODICAL:

Zhurnal Obshchey Khimii, 1957, vol 27, No 2, pp. 322-325 (U.S.S.R.)

ABSTRACT:

In order to explain the nature of the reaction between chloral and ether, the authors investigated the viscosity and density of systems: chloral - ethyl formate, chloral - ethyl acetate, chloral-ethyl ether, chloral - anisole and chloral - acetoacetic ester at temperatures ranging from 25 to 75°. Viscosity was found to be constant in the first four systems and in the fifth one it varies with time. A reaction between the components was established in all systems investigated but only the reaction of the chloral-acetoacetic ester system is clearly expressed and this is explained by the presence in the molecule of the latter of a hydroxyl group.

It was established that acetoacetic ester in the presence of pyridine reacts with chloral forming a compound which during distillation in vacuum

Card 1/2

Viscosity of Binary Systems with Chloral. Part 5 79-2-9/58

decomposes into chloral and acetoacetic ester. Optical investigation of acetoacetic ester solutions in pyridine showed that the latter like alcohol, reacts with the pyridine by means of the hydroxyl group. Also the reaction between the chloral and the acetoacetic acid is due to this hydroxyl group of enol form.

4 tables, 1 graph. There are 2 references, of which 2 are Slavic.

ASSOCIATION:

The Kiev Polytechnicum

PRESENTED BY:

SUBMITTED:

January 25, 1956

AVAILABLE:

Library of Congress

Card 2/2

~~UDOVENKO, H. V.~~: KHOMENKO, R. I.

Viscosity of binary systems containing chloral. Part 6. Zhru. ob. khim.
27 no.3:583-585 Mr '57. (MIRA 10:6)

1. Sredneaziatskiy gosudarstvennyy universitet i Kievskiy politekhnicheskii institut.
(Systems (Chemistry)) (Chloral)

UDOVENKO, V. V.

USSR/Inorganic Chemistry. Complex Compounds.

C

Abs Jour: Ref. Zhur-Khimiya, No 1, 1958, 677.

Author : Udovenko, V.V., Pialkov, Yu. Ya.

Inst : *Kiev Polytech Inst.*

Title : Interaction of Silicon Tetrachloride with Acetic and Benzoic Anhydrides.

Orig Pub: Zh. Obshch. Khimii, 1957, 27, No 4, 905-906.

Abstract: It was established that SiCl_4 can interact directly with anhydrides of organic acids thereby forming mixed anhydrides of organic and orthosilicic acids. This reaction proceeds under less rigid conditions than the reactions of interaction of SiCl_4 with corresponding acids. By means of the above reaction the tetracetate and tetrabenzoate of silicon were obtained in good yields.

Card : 1/1

-17-

AUTHORS: Udovenko, V. V., Fialkov, Yu. Ya. 79-28 -3-54/61

TITLE: The Viscosity of Binary Systems With a Substitution Reaction (Vyazkost' dvoynykh sistem s obmennym vzaimodeystviyem)

PERIODICAL: Zhurnal Obshchey Khimii, 1958, Vol. 28, Nr 3, pp. 814-818 (USSR)

ABSTRACT: The viscosity diagrams of binary systems are to a great extent systematized at present and are often used for the explanation of the occurring processes. When systems with chemical reactions are considered the present classification (Ref. 1) provides diagram types for such systems in which chemical reactions take place with a decrease of the molecular number, e. g. in the system water-chloral, or for systems in which the molecular number does not change, e. g. in the system acetic anhydride-water. Such systems have been little investigated. Therefore N. A. Trifonov suggested such model systems as, among other, diethyl-water, systems conducive to visualization of the type of the diagram of viscosity when only

Card 1/3

The Viscosity of Binary Systems With a Substitution
Reaction

79-28 -3-54/61

one chemical compound without a decrease of the molecular number is formed. It must be noted that in the system acetic anhydride-water the number of molecules remains the same after the chemical reaction had taken place as two molecules of acetic acid are formed from the molecules of this anhydride and water. The reaction leading to the formation of a chemical compound is not the only possible for reactions where the molecular number remains unchanged. Not less frequent is the substitution reaction where the final products are two chemical compounds. Systems of this kind are of great interest for the theory of physical and chemical analysis; they have, however, not been investigated by any chemical scientist with respect to the viscosity method. Below, data on the viscosity of systems are mentioned in which one component is silicon tetrachloride and the other one of the following compounds: methylal (dimethoxymethane), acetal (1,1-diethoxyethane) and acetic anhydride. According to the methods of viscosity and density the binary systems silicon tetrachloride-methylal at 20 and 30°C, silicon tetrachloride-acetal at 20, 30 and 40°C and silicon tetrachloride-acetic acid anhydride in benzene

Card 2/3

The Viscosity of Binary Systems With a Substitution Reaction 79-28 5-54/61

as indifferent medium at 20°C were investigated. The authors found that in the reaction of silicontetrachloride with methylal the final products are: dichlorodimethoxysilane and chlorodimethylether, and with acetal: dichloroethoxysilane and chlorodiethylether. This reaction represents a new method of the synthesis of dichlorodialkoxysilane which differs from the existing ones by its good yield and by the purity of the products. There are 1 figure, 3 tables and 5 references; 4 of which are Soviet.

ASSOCIATION: Kiyevskiy politekhnicheskii institut (Kiyev Polytechnical Institute)

SUBMITTED: March 3, 1957.

Card 3/3

AUTHORS: Udovenko, V. V., Aleksandrova, L. P. SOV/76-32-8-26/37

TITLE: The Solubility in the System Formic Acid - 1,2-Dichloro Ethane - Water (Rastvorimost' v sisteme murav'inaya kislota - 1,2-dikhloretan - voda)

PERIODICAL: Zhurnal fizicheskoy khimii, 1958, Vol. 32, Nr 8, pp. 1889-1892 (USSR)

ABSTRACT: The mutual solubility in the above mentioned ternary system was investigated within the temperature range of from 20 to 70° according to the polythermal method by V. F. Alekseyev. The data of the mutual solubility of water and dichloro ethane were taken from other publications. In the determinations of the solubility eight polythermal lines were taken at different constant ratios between formic acid and dichloro ethane, and the results obtained were given in a table. There exists a greater laminated section in the system which decreases on an increase in temperature. For determining the intersections in the system the compositions of the layers of equilibrium were determined by a titration of the acid in both layers at 30, 45 and 60°C. The kinetic points of the mutual solubility on the curves of the separation of layers were determined according

Card 1/2

SOV/76-32-8-26/37

The Solubility in the System Formic Acid - 1,2-Dichloro Ethane - Water

to the method mentioned above. The coefficients of distribution were calculated. The difference observed between the system $H_2O-C_2H_5OH-C_2H_4Cl_2$ and that investigated was explained by the greater coefficient of distribution of formic acid as compared to that of ethanol. There are 5 tables and 3 references, 2 of which are Soviet.

ASSOCIATION: Kiyevskiy politekhnicheskij institut (Kiyev Polytechnical Institute)

SUBMITTED: March 25, 1957

Card 2/2

5(4)

SOV/78-4-2-17/40

AUTHORS: Udovenko, V. V., Artemenko, M. V.

TITLE: On the Interaction of Copper Chloride With Monoethanolamine
(O vzaimodeystvii khlornoy medi s monoetanolaminom)

PERIODICAL: Zhurnal neorganicheskoy khimii, 1959, Vol 4, Nr 2,
pp 352-355 (USSR)

ABSTRACT: On the interaction of monoethanolamine with copper salts complex compounds of various compositions are formed. The following compounds are formed by copper chloride with ethanol amine combined with hydrochloric acid: $[\text{CH}_2\text{OHCH}_2\text{NH}_3]\text{CuCl}_3$ and $(\text{CH}_2\text{OHCH}_2\text{NH}_3)_2\cdot\text{CuCl}_4$. On the interaction of a concentrated alcoholic solution of copper chloride with monoethanol amine the compounds $\text{CH}_2\text{NH}_2\text{CH}_2\text{OCuCl}$ and $(\text{CH}_2\text{OHCH}_2\text{NH}_3)\text{CuCl}_3$ are formed. In weak solutions of copper chloride in acetone the compound $\text{Cu}(\text{CH}_2\text{OHCH}_2\text{NH}_2)_4\text{Cl}_2$ is formed. This compound crystallizes in well formed blue crystals whose melting point is 103° ; it is soluble in alcohol and water, and insoluble in acetone. On the interaction of copper chloride with $\text{Cu}(\text{CH}_2\text{OHCH}_2\text{NH}_2)_4\text{Cl}_2$

Card 1/2